

65

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
21 April 2005 (21.04.2005)

PCT

(10) International Publication Number  
**WO 2005/035525 A2**

(51) International Patent Classification<sup>7</sup>: C07D 403/00

(74) Agent: **HALEY, James, F., Jr.**; Fish & Neave, 1251 Avenue of the Americas, New York, NY 10020 (US).

(21) International Application Number:  
PCT/US2004/029093

(22) International Filing Date:  
7 September 2004 (07.09.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/500,670 5 September 2003 (05.09.2003) US

(71) Applicant (for all designated States except US): **VERTEX PHARMACEUTICALS INCORPORATED** [US/US]; 130 Waverly Street, Cambridge, MA 02139-4242 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **PERNI, Robert, B.** [US/US]; 130 Robert Road, Marlborough, MA 01752 (US). **COURT, John, J.** [US/US]; 27 Ernies Drive, Littleton, MA 01460 (US). **BRITT, Shawn, D.** [US/US]; 56 Marilyn Road, Andover, MA 01810 (US). **PITLIK, Janos** [HU/US]; 1 Robin Circle, Westborough, MA 01581 (US). **VAN DRIE, John, H.** [US/US]; 34 Stinson Road, Andover, MA 01810 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **INHIBITORS OF SERINE PROTEASES, PARTICULARLY HCV NS3-NS4A PROTEASE**

(57) Abstract: The present invention relates to compounds that inhibit serine protease activity, particularly the activity of hepatitis C virus NS3-NS4A protease. As such, they act by interfering with the life cycle of the hepatitis C virus and are also useful as antiviral agents. The invention further relates to compositions comprising these compounds either for *ex vivo* use for administration to a patient suffering from HCV infection. The invention also relates to methods of treating an HCV infection in a patient by administering a composition comprising a compound of this invention. The invention further relates to processes for preparing these compounds.

WO 2005/035525 A2

INHIBITORS OF SERINE PROTEASES,  
PARTICULARLY HCV NS3-NS4A PROTEASE

5

TECHNICAL FIELD OF THE INVENTION

[0001] The present invention relates to compounds that  
10 inhibit serine protease activity, particularly the  
activity of hepatitis C virus NS3-NS4A protease. As  
such, they act by interfering with the life cycle of the  
hepatitis C virus and are also useful as antiviral  
agents. The invention also relates to processes for  
15 preparing these compounds. The invention further relates  
to compositions comprising these compounds either for ex  
vivo use or for administration to a patient suffering  
from HCV infection. The invention also relates to  
methods of treating an HCV infection in a patient by  
20 administering a composition comprising a compound of this  
invention.

BACKGROUND OF THE INVENTION

[0002] Infection by hepatitis C virus ("HCV") is a  
compelling human medical problem. HCV is recognized as  
25 the causative agent for most cases of non-A, non-B  
hepatitis, with an estimated human sero-prevalence of 3%  
globally [A. Alberti et al., "Natural History of  
Hepatitis C," J. Hepatology, 31., (Suppl. 1), pp. 17-24  
(1999)]. Nearly four million individuals may be infected  
30 in the United States alone [M.J. Alter et al., "The  
Epidemiology of Viral Hepatitis in the United States,  
Gastroenterol. Clin. North Am., 23, pp. 437-455 (1994);  
M. J. Alter "Hepatitis C Virus Infection in the United  
States," J. Hepatology, 31., (Suppl. 1), pp. 88-91  
35 (1999)].

[0003] Upon first exposure to HCV only about 20% of infected individuals develop acute clinical hepatitis while others appear to resolve the infection spontaneously. In almost 70% of instances, however, the virus establishes a chronic infection that persists for decades [S. Iwarson, "The Natural Course of Chronic Hepatitis," FEMS Microbiology Reviews, 14, pp. 201-204 (1994); D. Lavanchy, "Global Surveillance and Control of Hepatitis C," J. Viral Hepatitis, 6, pp. 35-47 (1999)]. This usually results in recurrent and progressively worsening liver inflammation, which often leads to more severe disease states such as cirrhosis and hepatocellular carcinoma [M.C. Kew, "Hepatitis C and Hepatocellular Carcinoma", FEMS Microbiology Reviews, 14, pp. 211-220 (1994); I. Saito et. al., "Hepatitis C Virus Infection is Associated with the Development of Hepatocellular Carcinoma," Proc. Natl. Acad. Sci. USA, 87, pp. 6547-6549 (1990)]. Unfortunately, there are no broadly effective treatments for the debilitating progression of chronic HCV.

[0004] The HCV genome encodes a polyprotein of 3010-3033 amino acids [Q.L. Choo, et. al., "Genetic Organization and Diversity of the Hepatitis C Virus." Proc. Natl. Acad. Sci. USA, 88, pp. 2451-2455 (1991); N. Kato et al., "Molecular Cloning of the Human Hepatitis C Virus Genome From Japanese Patients with Non-A, Non-B Hepatitis," Proc. Natl. Acad. Sci. USA, 87, pp. 9524-9528 (1990); A. Takamizawa et. al., "Structure and Organization of the Hepatitis C Virus Genome Isolated From Human Carriers," J. Virol., 65, pp. 1105-1113 (1991)]. The HCV nonstructural (NS) proteins are presumed to provide the essential catalytic machinery for viral replication. The NS proteins are derived by proteolytic cleavage of the polyprotein [R.

Bartenschlager et. al., "Nonstructural Protein 3 of the Hepatitis C Virus Encodes a Serine-Type Proteinase Required for Cleavage at the NS3/4 and NS4/5 Junctions," J. Virol., 67, pp. 3835-3844 (1993); A. Grakoui et. al.,  
5 "Characterization of the Hepatitis C Virus-Encoded Serine Proteinase: Determination of Proteinase-Dependent Polyprotein Cleavage Sites," J. Virol., 67, pp. 2832-2843 (1993); A. Grakoui et. al., "Expression and Identification of Hepatitis C Virus Polyprotein Cleavage  
10 Products," J. Virol., 67, pp. 1385-1395 (1993); L. Tomei et. al., "NS3 is a serine protease required for processing of hepatitis C virus polyprotein", J. Virol., 67, pp. 4017-4026 (1993)].

[0005] The HCV NS protein 3 (NS3) contains a serine  
15 protease activity that helps process the majority of the viral enzymes, and is thus considered essential for viral replication and infectivity. It is known that mutations in the yellow fever virus NS3 protease decreases viral infectivity [Chambers, T.J. et. al., "Evidence that the  
20 N-terminal Domain of Nonstructural Protein NS3 From Yellow Fever Virus is a Serine Protease Responsible for Site-Specific Cleavages in the Viral Polyprotein", Proc. Natl. Acad. Sci. USA, 87, pp. 8898-8902 (1990)]. The first 181 amino acids of NS3 (residues 1027-1207 of the  
25 viral polyprotein) have been shown to contain the serine protease domain of NS3 that processes all four downstream sites of the HCV polyprotein [C. Lin et al., "Hepatitis C Virus NS3 Serine Proteinase: *Trans*-Cleavage Requirements and Processing Kinetics", J. Virol., 68, pp.  
30 8147-8157 (1994)].

[0006] The HCV NS3 serine protease and its associated cofactor, NS4A, helps process all of the viral enzymes, and is thus considered essential for viral replication. This processing appears to be analogous to that carried

out by the human immunodeficiency virus aspartyl protease, which is also involved in viral enzyme processing HIV protease inhibitors, which inhibit viral protein processing are potent antiviral agents in man, indicating that interrupting this stage of the viral life cycle results in therapeutically active agents. Consequently it is an attractive target for drug discovery.

[0007] Several potential HCV protease inhibitors have been described [PCT publication Nos. WO 02/18369, WO 02/08244, WO 00/09558, WO 00/09543, WO 99/64442, WO 99/07733, WO 99/07734, WO 99/50230, WO 98/46630, WO 98/17679 and WO 97/43310, United States Patent 5,990,276, M. Llinas-Brunet et al., Bioorg. Med. Chem. Lett., 8, pp. 1713-18 (1998); W. Han et al., Bioorg. Med. Chem. Lett., 10, 711-13 (2000); R. Dunsdon et al., Bioorg. Med. Chem. Lett., 10, pp. 1571-79 (2000); M. Llinas-Brunet et al., Bioorg. Med. Chem. Lett., 10, pp. 2267-70 (2000); and S. LaPlante et al., Bioorg. Med. Chem. Lett., 10, pp. 2271-74 (2000)].


[0008] There are not currently any satisfactory anti-HCV agents or treatments. The only established therapy for HCV disease is interferon treatment. However, interferons have significant side effects [M. A. Wlaker et al., "Hepatitis C Virus: An Overview of Current Approaches and Progress," DDT, 4, pp. 518-29 (1999); D. Moradpour et al., "Current and Evolving Therapies for Hepatitis C," Eur. J. Gastroenterol. Hepatol., 11, pp. 1199-1202 (1999); H. L. A. Janssen et al. "Suicide Associated with Alfa-Interferon Therapy for Chronic Viral Hepatitis," J. Hepatol., 21, pp. 241-243 (1994); P.F. Renault et al., "Side Effects of Alpha Interferon," Seminars in Liver Disease, 9, pp. 273-277. (1989)] and induce long term remission in only a fraction (~ 25%) of

cases [O. Weiland, "Interferon Therapy in Chronic Hepatitis C Virus Infection" , FEMS Microbiol. Rev., 14, pp. 279-288 (1994)]. Moreover, the prospects for effective anti-HCV vaccines remain uncertain.

- 5 [0009] Thus, there is a need for compounds useful in anti-HCV therapies. Such compounds would have therapeutic potential as protease inhibitors, particularly as serine protease inhibitors, and more particularly as HCV NS3 protease inhibitors. These  
10 compounds would be useful as antiviral agents, particularly as anti-HCV agents. There is a particular need for compounds with improved enzyme inhibition or cellular activity.

#### SUMMARY OF THE INVENTION

- 15 [0010] The present invention addresses these needs by providing a compound of formula I:




I

wherein the variables are as defined herein.

- 20 [0011] The invention also relates to compositions that comprise the above compounds and the use thereof. Such compositions may be used to pre-treat invasive devices to be inserted into a patient, to treat biological samples, such as blood, prior to administration to a patient, and  
25 for direct administration to a patient. In each case the composition will be used to inhibit HCV replication and to lessen the risk of or the severity of HCV infection.
- [0012] The invention also relates to processes for preparing the compounds of formula I.

DETAILED DESCRIPTION OF THE INVENTION

[0013] The present invention provides a compound of formula I:



I

wherein:

Ar is a 5- to 10-membered aromatic ring having up to 4 heteroatoms selected from O, S, N(H), SO, and SO<sub>2</sub>, wherein 1 to 3 ring atoms are optionally and independently substituted with J;

R<sup>1</sup> and R<sup>2</sup> are independently:

(C1-C12)-aliphatic-,

(C3-C10)-cycloalkyl- or -cycloalkenyl-,

[(C3-C10)-cycloalkyl- or -cycloalkenyl]-(C1-C12)-

aliphatic-,

(C6-C10)-aryl-(C1-C12)aliphatic-,

(C6-C10)-heteroaryl-(C1-C12)aliphatic-,

wherein up to 3 aliphatic carbon atoms in R<sup>1</sup> and R<sup>2</sup> may be replaced by a heteroatom selected from O, N, S, SO, or SO<sub>2</sub> in a chemically stable arrangement;

wherein each of R<sup>1</sup> and R<sup>2</sup> is independently and optionally substituted with up to 3 substituents independently selected from J;

R<sup>3</sup> and R<sup>3'</sup> are independently hydrogen or (C1-C12)-

aliphatic, wherein any hydrogen is optionally replaced with halogen; wherein any terminal carbon atom of R<sup>3</sup> is optionally substituted with sulfhydryl or hydroxy; or R<sup>3</sup> is phenyl or -CH<sub>2</sub>phenyl, wherein said phenyl group is optionally substituted with up to 3 substituents

independently selected from J; or

$R^3$  and  $R^{3'}$  together with the atom to which they are bound is a 3- to 6-membered ring having up to 2 heteroatoms selected from N, NH, O, SO, or  $SO_2$ ; wherein the ring has up to 2 substituents selected independently from J;

5  $R^4$  and  $R^{4'}$  are independently:

hydrogen-,

(C1-C12)-aliphatic-,

(C3-C10)-cycloalkyl- or -cycloalkenyl-,

(C3-C10)-cycloalkyl-(C1-C12)-aliphatic-,

10 (C6-C10)-aryl-,

(C3-C10)-heterocyclyl-; or



(C5-C10)-heteroaryl-;

wherein up to two aliphatic carbon atoms in  $R^4$  and  $R^{4'}$  may be replaced by a heteroatom selected from O, N,

15 S, SO, or  $SO_2$ ;

wherein each of  $R^4$  and  $R^{4'}$  is independently and optionally substituted with up to 3 substituents independently selected from J;

W is:



wherein each  $R_6$  is independently:

hydrogen-,

(C1-C12)-aliphatic-,

(C6-C10)-aryl-,

25 (C6-C10)-aryl-(C1-C12)aliphatic-,

(C3-C10)-cycloalkyl- or cycloalkenyl-,



[(C3-C10)-cycloalkyl- or cycloalkenyl]-(C1-C12)-  
aliphatic-,

(C3-C10)-heterocyclyl-,

(C3-C10)-heterocyclyl-(C1-C12)-aliphatic-,

5 (C5-C10)heteroaryl-, or

(C5-C10)heteroaryl-(C1-C12)-aliphatic-, or

two R<sub>6</sub> groups, which are bound to the same  
nitrogen atom, form together with that nitrogen  
atom, a (C3-C10)-heterocyclic ring;

10 wherein R<sub>6</sub> is optionally substituted with up to 3  
J substituents;

wherein each R<sub>8</sub> is independently -OR'; or the R<sub>8</sub>  
groups together with the boron atom, is a (C3-C10)-  
membered heterocyclic ring having in addition to the  
15 boron up to 3 additional heteroatoms selected from N,  
NH, O, SO, and SO<sub>2</sub>;

T is:

(C1-C12)-aliphatic-;

(C6-C10)-aryl-,

20 (C6-C10)-aryl-(C1-C12)aliphatic-,

(C5-C10)heteroaryl-, or

(C5-C10)heteroaryl-(C1-C12)-aliphatic-;

wherein each T is optionally substituted with up to  
3 J substituents;

25 J is halogen, -OR', -NO<sub>2</sub>, -CN, -CF<sub>3</sub>, -OCF<sub>3</sub>, -R', oxo,  
thioxo, 1,2-methylenedioxy, 1,2-ethylenedioxy, =N(R'),  
=N(OR'), -N(R')<sub>2</sub>, -SR', -SOR', -SO<sub>2</sub>R', -SO<sub>2</sub>N(R')<sub>2</sub>,  
-SO<sub>3</sub>R', -C(O)R', -C(O)C(O)R', -C(O)CH<sub>2</sub>C(O)R', -C(S)R',  
-C(S)OR', -C(O)OR', -C(O)C(O)OR', -C(O)C(O)N(R')<sub>2</sub>,  
30 -OC(O)R', -C(O)N(R')<sub>2</sub>, -OC(O)N(R')<sub>2</sub>, -C(S)N(R')<sub>2</sub>,  
-(CH<sub>2</sub>)<sub>0-2</sub>NHC(O)R', -N(R')N(R')COR', -N(R')N(R')C(O)OR',  
-N(R')N(R')CON(R')<sub>2</sub>, -N(R')SO<sub>2</sub>R', -N(R')SO<sub>2</sub>N(R')<sub>2</sub>,  
-N(R')C(O)OR', -N(R')C(O)R', -N(R')C(S)R',  
-N(R')C(O)N(R')<sub>2</sub>, -N(R')C(S)N(R')<sub>2</sub>, -N(COR')COR',

-N(OR')R', -C(=NH)N(R')<sub>2</sub>, -C(O)N(OR')R', -C(=NOR')R',  
-OP(O)(OR')<sub>2</sub>, -P(O)(R')<sub>2</sub>, -P(O)(OR')<sub>2</sub>, or -P(O)(H)(OR');


R' is independently selected from:

- hydrogen-,  
5 (C1-C12)-aliphatic-,  
(C3-C10)-cycloalkyl- or -cycloalkenyl-,  
[(C3-C10)-cycloalkyl or -cycloalkenyl]-(C1-C12)-  
aliphatic-,  
(C6-C10)-aryl-,  
10 (C6-C10)-aryl-(C1-C12)aliphatic-,  
(C3-C10)-heterocyclyl-,  
(C6-C10)-heterocyclyl-(C1-C12)aliphatic-,  
(C5-C10)-heteroaryl-, or  
(C5-C10)-heteroaryl-(C1-C12)-aliphatic-;  
15 wherein R' is optionally substituted with up to 3  
J groups;

wherein two R' groups bound to the same atom  
form a 3- to 10-membered aromatic or non-aromatic  
ring having up to 3 heteroatoms independently  
20 selected from N, O, S, SO, or SO<sub>2</sub>, wherein said  
ring is optionally fused to a (C6-C10)aryl, (C5-  
C10)heteroaryl, (C3-C10)cycloalkyl, or a (C3-  
C10)heterocyclyl, wherein any ring has up to 3  
substituents selected independently from J.


25 [0014] These compounds solve the above problems by  
providing compounds with improved enzyme and/or cell  
activity. For example, compounds of this invention,  
particularly the preferred compounds exhibit better  
enzyme inhibition than the compounds of WO 98/17679. The  
30 compounds of this invention also have better cellular  
data than other reported compounds (see the documents  
cited herein).

[0015] In one embodiment, if  $R^1$  is cyclohexyl,  $R^2$  is *t*-butyl,  $R^{3'}$  is H,  $R^3$  is *n*-propyl, W is  $-\text{C}(\text{O})\text{C}(\text{O})\text{N}(\text{H})-$ cyclopropyl, and T is:



5 ; then Ar is 4-quinazoline or 5-chloro-2-pyridyl.

[0016] The present invention also provides a compound of formula I:



I

10 wherein:

$n$  is 0 or 1;

Ar is a 5- to 10-membered aromatic ring having up to 4 heteroatoms selected from O, S, N(H), SO, and SO<sub>2</sub>, wherein 1 to 3 ring atoms are optionally and

15 independently substituted with J;

$R^1$ ,  $R^2$ ,  $R^{12}$ , and  $R^{13}$  are independently:

(C1-C12)-aliphatic-,

(C3-C10)-cycloalkyl- or -cycloalkenyl-,

[(C3-C10)-cycloalkyl- or -cycloalkenyl]-(C1-C12)-

20 aliphatic-,

(C6-C10)-aryl-(C1-C12)aliphatic-,

(C6-C10)-heteroaryl-(C1-C12)aliphatic-,


wherein up to 3 aliphatic carbon atoms in  $R^1$  and  $R^2$  may be replaced by a heteroatom selected from O, N, NH, S, SO, or SO<sub>2</sub> in a chemically stable arrangement;

25 wherein each of  $R^1$  and  $R^2$  is independently and optionally substituted at each substitutable position

with up to 3 substituents independently selected from J;

- 5  $R^3$  and  $R^{3'}$  are independently hydrogen or (C1-C12)-aliphatic, wherein any hydrogen is optionally replaced with halogen; wherein any terminal carbon atom of  $R^3$  is optionally substituted with sulfhydryl or hydroxy; or  $R^3$  is phenyl or  $-\text{CH}_2\text{phenyl}$ , wherein said phenyl group is optionally substituted with up to 3 substituents independently selected from J; or
- 10  $R^3$  and  $R^{3'}$  together with the atom to which they are bound is a 3- to 6-membered ring having up to 2 heteroatoms selected from N, NH, O, SO, and  $\text{SO}_2$ ; wherein the ring has up to 2 substituents selected independently from J;
- 15  $R^4$  and  $R^{4'}$  are independently:  
hydrogen-,  
(C1-C12)-aliphatic-,  
(C3-C10)-cycloalkyl- or -cycloalkenyl-,  
(C3-C10)-cycloalkyl-(C1-C12)-aliphatic-,  
20 (C6-C10)-aryl-,  
(C3-C10)-heterocyclyl-; or  
(C5-C10)-heteroaryl-;  
wherein up to two aliphatic carbon atoms in  $R^4$  and  $R^{4'}$  may be replaced by a heteroatom selected from O, N,  
25 S, SO, and  $\text{SO}_2$ ;  
wherein each of  $R^4$  and  $R^{4'}$  is independently and optionally substituted with up to 3 substituents independently selected from J;

W is:



wherein

$Y$  is  $-CO_2H$ , a derivative of  $-CO_2H$ , or a bioisostere of  $-CO_2H$ ;

- 5 each  $R_6$  is independently:
- hydrogen-,
  - (C1-C12)-aliphatic-,
  - (C6-C10)-aryl-,
  - (C6-C10)-aryl-(C1-C12)aliphatic-,
  - 10 (C3-C10)-cycloalkyl- or cycloalkenyl-,
  - [(C3-C10)-cycloalkyl- or cycloalkenyl]-(C1-C12)-
  - aliphatic-,
  - (C3-C10)-heterocyclyl-,
  - (C3-C10)-heterocyclyl-(C1-C12)-aliphatic-,
  - 15 (C5-C10)heteroaryl-, or
  - (C5-C10)heteroaryl-(C1-C12)-aliphatic-, or
  - two  $R_6$  groups, which are bound to the same
  - nitrogen atom, form together with that nitrogen
  - atom, a (C3-C10)-heterocyclic ring;
  - 20 wherein  $R_6$  is optionally substituted with up to 3
  - $J$  substituents;

wherein each  $R_8$  is independently  $-OR'$ ; or the  $R_8$  groups together with the boron atom, is a (C3-C10)-membered heterocyclic ring having in addition to the

boron up to 3 additional heteroatoms selected from N, NH, O, SO, and SO<sub>2</sub>;

T is:

(C1-C12)-aliphatic-;

5 (C6-C10)-aryl-,

(C6-C10)-aryl-(C1-C12)aliphatic-,

(C5-C10)heteroaryl-, or

(C5-C10)heteroaryl-(C1-C12)-aliphatic-;

10 wherein each T is optionally substituted with up to 3 J substituents; and

wherein up to 3 aliphatic carbon atoms in T may be replaced by a heteroatom selected from O, N, NH, S, SO, or SO<sub>2</sub> in a chemically stable arrangement;

provided that if T is pyrrole, the pyrrole is not substituted at the 3-position with J, with J being -C(O)R', -C(O)C(O)R', -C(O)CH<sub>2</sub>C(O)R', -C(S)R', -C(S)OR', -C(O)OR', -C(O)C(O)OR', -C(O)C(O)N(R')<sub>2</sub>, -C(O)N(R')<sub>2</sub>, -C(S)N(R')<sub>2</sub>, -C(=NH)N(R')<sub>2</sub>, -C(O)N(OR')R', or -C(=NOR')R';

J is halogen, -OR', -NO<sub>2</sub>, -CN, -CF<sub>3</sub>, -OCF<sub>3</sub>, -R', oxo, thioxo, 1,2-methylenedioxy, 1,2-ethylenedioxy, =N(R'), =N(OR'), -N(R')<sub>2</sub>, -SR', -SOR', -SO<sub>2</sub>R', -SO<sub>2</sub>N(R')<sub>2</sub>, -SO<sub>3</sub>R', -C(O)R', -C(O)C(O)R', -C(O)CH<sub>2</sub>C(O)R', -C(S)R', -C(S)OR', -C(O)OR', -C(O)C(O)OR', -C(O)C(O)N(R')<sub>2</sub>, -OC(O)R', -C(O)N(R')<sub>2</sub>, -OC(O)N(R')<sub>2</sub>, -C(S)N(R')<sub>2</sub>, - (CH<sub>2</sub>)<sub>0-2</sub>NHC(O)R', -N(R')N(R')COR', -N(R')N(R')C(O)OR', -N(R')N(R')CON(R')<sub>2</sub>, -N(R')SO<sub>2</sub>R', -N(R')SO<sub>2</sub>N(R')<sub>2</sub>, -N(R')C(O)OR', -N(R')C(O)R', -N(R')C(S)R', -N(R')C(O)N(R')<sub>2</sub>, -N(R')C(S)N(R')<sub>2</sub>, -N(COR')COR', -N(OR')R', -C(=NH)N(R')<sub>2</sub>, -C(O)N(OR')R', -C(=NOR')R', -OP(O)(OR')<sub>2</sub>, -P(O)(R')<sub>2</sub>, -P(O)(OR')<sub>2</sub>, or -P(O)(H)(OR');

R' is:

hydrogen-,

(C1-C12)-aliphatic-,

(C3-C10)-cycloalkyl- or -cycloalkenyl-,

[(C3-C10)-cycloalkyl or -cycloalkenyl]-(C1-C12)-  
aliphatic-,

(C6-C10)-aryl-,

(C6-C10)-aryl-(C1-C12)aliphatic-,

5 (C3-C10)-heterocyclyl-,

(C6-C10)-heterocyclyl-(C1-C12)aliphatic-,

(C5-C10)-heteroaryl-, or


(C5-C10)-heteroaryl-(C1-C12)-aliphatic-;

wherein R' is optionally substituted with up to 3

10 J groups;

wherein two R' groups bound to the same atom form a 3- to  
10-membered aromatic or non-aromatic ring having up to 3  
heteroatoms independently selected from N, O, S, SO, or  
SO<sub>2</sub>, wherein said ring is optionally fused to a (C6-  
15 C10)aryl, (C5-C10)heteroaryl, (C3-C10)cycloalkyl, or a  
(C3-C10)heterocyclyl, wherein any ring has up to 3  
substituents selected independently from J.

[0017] In one embodiment, if R<sup>1</sup> is cyclohexyl, R<sup>2</sup> is t-  
butyl, R<sup>3'</sup> is H, R<sup>3</sup> is n-propyl, W is -C(O)C(O)N(H)-  
20 cyclopropyl, and T is:




; then Ar is not 4-quinazoline or 5-chloro-2-  
pyridyl.

[0018] In another embodiment, if T is pyrrole, the  
pyrrole is not substituted with J, but is optionally  
25 fused to a 5-membered or a 6-membered aryl or heteroaryl  
ring.


[0019] In another embodiment, if T is pyrrole, the  
pyrrole is not substituted at the 3-position with J, but  
is optionally fused to a 5-membered or a 6-membered aryl  
30 or heteroaryl ring.

[0020] In another embodiment, T is not




[0021] In another embodiment, T is not pyrrole.

[0022] In a more specific embodiment, this invention  
 5 provides a compound wherein n is 0 and the compound has  
 the formula I-A:




I-A;

wherein the variable are as defined in any of the  
 10 embodiments herein. In a compound of formula A, W is  
 preferably:




[0023] In another specific embodiment, this invention  
 provides a compound wherein n is 1 and the compound has  
 15 the formula I-B:



I-B;




wherein the variable are as defined in any of the embodiments herein. In a compound of formula I-B, W is preferably:




## 5 Definitions

[0024] The term "aryl" as used herein means a monocyclic or bicyclic carbocyclic aromatic ring system. Phenyl is an example of a monocyclic aromatic ring system. Bicyclic aromatic ring systems include systems wherein both rings are aromatic, e.g., naphthyl, and systems wherein only one of the two rings is aromatic, e.g., tetralin.

[0025] The term "bioisostere"  $-CO_2H$  as used in herein refers to a chemical moiety which may substitute for a carboxylic acid group in a biologically active molecule. Examples of such groups are disclosed in Christopher A. Lipinski, "Bioisosteres in Drug Design" Annual Reports in Medicinal Chemistry, 21, pp. 286-88 (1986), and in C. W. Thornber, "Isosterism and Molecular Modification in Drug Design" Chemical Society Reviews, pp. 563-580 (1979). Examples of such groups include, but are not limited to,  $-COCH_2OH$ ,  $-CONHOH$ ,  $SO_2NHR'$ ,  $-SO_3H$ ,  $-PO(OH)NH_2$ ,  $-CONHCN$ ,  $-OSO_3H$ ,  $-CONHSO_2R'$ ,  $-PO(OH)_2$ ,  $-PO(OH)(OR')$ ,  $-PO(OH)(R')$ ,  $-OPO(OH)_2$ ,  $-OPO(OH)(OR')$ ,  $-OPO(OH)(R')$ ,  $HNPO(OH)_2$ ,  $-NHPO(OH)(OR')$ ,  $-NHPO(OH)(R')$





5        [0026]        The term "heterocyclyl" as used herein means a  
monocyclic or bicyclic non-aromatic ring system having 1  
to 3 heteroatom or heteroatom groups in each ring  
selected from O, N, NH, S, SO, or SO<sub>2</sub> in a chemically  
stable arrangement. In a bicyclic non-aromatic ring  
system embodiment of "heterocyclyl" one or both rings may  
10        contain said heteroatom or heteroatom groups.

[0027] Examples of heterocyclic rings include 3-1H-benzimidazol-2-one, 3-(1-alkyl)-benzimidazol-2-one, 2-tetrahydrofuran-2-yl, 3-tetrahydrofuran-2-yl, 2-tetrahydrothiophen-2-yl, 3-tetrahydrothiophen-2-yl, 2-morpholino, 3-morpholino, 4-morpholino, 2-thiomorpholino, 3-thiomorpholino, 4-thiomorpholino, 1-pyrrolidin-2-yl, 2-pyrrolidin-3-yl, 3-pyrrolidin-1-yl, 1-tetrahydropiperazin-2-yl, 2-tetrahydropiperazin-3-yl, 3-tetrahydropiperazin-1-yl, 1-piperidin-2-yl, 2-piperidin-3-yl, 3-piperidin-1-yl, 1-pyrazolin-3-yl, 3-pyrazolin-4-yl, 5-pyrazolin-1-yl, 1-piperidin-2-yl, 2-piperidin-3-yl, 3-piperidin-4-yl, 2-thiazolidin-3-yl, 3-thiazolidin-4-yl, 4-thiazolidin-1-yl,

imidazolidinyl, 2-imidazolidinyl, 4-imidazolidinyl, 5-imidazolidinyl, indolinyl, tetrahydroquinolinyl, tetrahydroisoquinolinyl, benzothiolane, benzodithiane, and 1,3-dihydro-imidazol-2-one.

5    **[0028]**    The term "heteroaryl" as used herein means a monocyclic or bicyclic aromatic ring system having 1 to 3 heteroatom or heteroatom groups in each ring selected from O, N, NH or S in a chemically stable arrangement. In such a bicyclic aromatic ring system embodiment of  
10    "heteroaryl":

- one or both rings may be aromatic; and
- one or both rings may contain said heteroatom or heteroatom groups.

**[0029]**    Examples of heteroaryl rings include 2-furanyl,  
15    3-furanyl, N-imidazolyl, 2-imidazolyl, 4-imidazolyl, 5-imidazolyl, benzimidazolyl, 3-isoxazolyl, 4-isoxazolyl, 5-isoxazolyl, 2-oxazolyl, 4-oxazolyl, 5-oxazolyl, N-pyrrolyl, 2-pyrrolyl, 3-pyrrolyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-pyrimidinyl, 4-pyrimidinyl, 5-pyrimidinyl,  
20    pyridazinyl (e.g., 3-pyridazinyl), 2-thiazolyl, 4-thiazolyl, 5-thiazolyl, tetrazolyl (e.g., 5-tetrazolyl), triazolyl (e.g., 2-triazolyl and 5-triazolyl), 2-thienyl, 3-thienyl, benzofuryl, benzothiophenyl, indolyl (e.g., 2-indolyl), pyrazolyl (e.g., 2-pyrazolyl), isothiazolyl,  
25    1,2,3-oxadiazolyl, 1,2,5-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,3-triazolyl, 1,2,3-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,5-thiadiazolyl, purinyl, pyrazinyl, 1,3,5-triazinyl, quinolinyl (e.g., 2-quinolinyl, 3-quinolinyl, 4-quinolinyl), and isoquinolinyl (e.g., 1-isoquinolinyl, 3-  
30    isoquinolinyl, or 4-isoquinolinyl).

**[0030]**    The term "aliphatic" as used herein means a straight chained or branched alkyl, alkenyl or alkynyl. It is understood that alkenyl or alkynyl embodiments need at least two carbon atoms in the aliphatic chain.

The term "cycloalkyl or cycloalkenyl" refers to a monocyclic or fused or bridged bicyclic carbocyclic ring system that is not aromatic. Cycloalkenyl rings have one or more units of unsaturation. Preferred cycloalkyl groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclohexenyl, cycloheptyl, cycloheptenyl, nornbornyl, adamantyl and decalin-yl.


[0031] As used herein, the carbon atom designations may have the indicated integer and any intervening integer. For example, the number of carbon atoms in a (C1-C4)-alkyl group is 1, 2, 3, or 4. It should be understood that these designation refer to the total number of atoms in the appropriate group. For example, in a (C3-C10)-heterocyclyl the total number of carbon atoms and heteroatoms is 3 (as in aziridine), 4, 5, 6 (as in morpholine), 7, 8, 9, or 10.




[0032] The phrase "chemically stable arrangement" as used herein refers to a compound structure that renders the compound sufficiently stable to allow manufacture and administration to a mammal by methods known in the art. Typically, such compounds are stable at a temperature of 40°C or less, in the absence of moisture or other chemically reactive condition, for at least a week.

#### Preferred Embodiments


[0033] In another embodiment of this invention, Ar is phenyl, pyridyl, quinolinyl, pyrimidinyl, or naphthyl, wherein each group is optionally substituted with 1, 2, or 3 J groups.

[0034] In another embodiment of this invention, Ar is






10 [0035] In another embodiment of this invention, Ar is:



[0036] In a particularly favorable embodiment, Ar is:



15

[0037] In another embodiment of this invention, Ar is a 6 or a 10-membered aromatic ring having 0, 1, or 2 nitrogen heteroatoms, wherein 1, 2, or 3 ring atoms are optionally and independently substituted with J.

20 [0038] In any of the embodiments of this invention, each J group on Ar is independently OR', NO<sub>2</sub>, CN, CF<sub>3</sub>, OCF<sub>3</sub>, R', COR', C(O)OR', C(O)N(R')<sub>2</sub>, SO<sub>2</sub>R', SO<sub>2</sub>N(R')<sub>2</sub>, 1,2-methylenedioxy, 1,2-ethylene dioxy, or NR'C(O)OR', NR'SO<sub>2</sub>R'.

25 [0039] In any of embodiments of this invention, each J group on Ar is preferably, and independently, OR',


halogen, CN, CF<sub>3</sub>, R', or COR'. More preferably, this J is halogen (particularly chloro).

[0040] In another embodiment of this invention, each J group on Ar is independently halo, trifluoromethyl, methyl, or NO<sub>2</sub>.

[0041] According to any preferred embodiment, T is (C<sub>6</sub>-C<sub>10</sub>)-aryl- or (C<sub>5</sub>-C<sub>10</sub>)heteroaryl-, wherein each T is optionally substituted with 1, 2, or 3 J substituents.


[0042] In a preferred embodiment, T is a 6-membered or a 10-membered aryl group. In another preferred embodiment, T is a 6-membered heteroaryl group that is optionally fused to another 5- or 6-membered aryl or heteroaryl group.

[0043] More preferred embodiments are those wherein T:



15


[0044] In a more preferred embodiment, T is:



or pyrazine.


[0045] In certain embodiments of this invention, any T group is optionally fused to a 5-membered or a 6-membered aryl or heteroaryl group.

5 [0046] Accordingly, one embodiment of this invention provides compounds wherein T is:



wherein each T group is optionally fused to a 5-membered or a 6-membered aryl or heteroaryl group.

10 [0047] In another embodiment, T is:



wherein T is optionally fused to a 5-membered or a 6-membered aryl or heteroaryl group.

[0048] In certain embodiments, T is not fused to  
15 another ring.


[0049] In any embodiment of this invention, each aryl or heteroaryl group of T is optionally and independently substituted with 1, 2, or 3 groups selected from -CH<sub>3</sub>, -H<sub>2</sub>CH<sub>3</sub>, halogen, acetyl, -CO<sub>2</sub>H, -(C1-C6-alkyl)-CO<sub>2</sub>H, or  
20 -O<sub>2</sub>R'.

[0050] In certain embodiments, 1 aliphatic carbon atom in T is replaced by a heteroatom selected from O, N, NH, S, SO, or SO<sub>2</sub> in a chemically stable arrangement. The heteroatom is preferably, O or NH. In an alternative  
5 embodiment no aliphatic carbon atom is replaced with T.

[0051] In certain compounds of this invention, an aliphatic group links to T to the remainder of the molecule. In general, in these aliphatic linkers replacement of an aliphatic carbon atom is preferred in  
10 compounds wherein n is 0. In compounds wherein n is 1, this aliphatic linker group, preferably, has no heteroatom replacements. Furthermore, compounds wherein n is 1, preferably, have no aliphatic linker group.

[0052] According to a preferred embodiment, W is  
15 -C(O)-C(O)-R<sub>6</sub>. Preferably, R<sub>6</sub> is phenyl, pyridyl, (C3-C6)-alkyl, (C3-C6)-cycloalkyl, -OH, -O-(C1-C6)-alkyl, -N(H)-(C3-C6)-cycloalkyl, -N(H)-C(H)(CH<sub>3</sub>)-(C6-C10)aryl, -N(H)-C(H)(CH<sub>3</sub>)-(C3-C10)-heterocyclyl, or -N(H)-C(H)(CH<sub>3</sub>)-(C5-C10)-heteroaryl, wherein each aryl, heterocyclyl, and  
20 heteroaryl is optionally substituted with halogen.

Preferred embodiments are selected from:



More preferably, R<sub>6</sub> is isopropyl.


[0053] According to one embodiment, W is -C(O)-C(O)-  
25 OR<sub>6</sub>. In this embodiment, R<sub>6</sub> is preferably hydrogen, (C1-C12)-aliphatic (more preferably, C1-C6-alkyl), (C6-C10)-aryl, (C3-C10)-cycloalkyl or -cycloalkenyl, (C3-C10)-



heterocyclyl, (C5-C10)heteroaryl, or C3-C6-cycloalkyl-(C1-C3)-alkyl, wherein the cycloalkyl is preferably a cyclopropyl group. The aryl group is optionally substituted with up to 3 J groups, wherein J is halogen, preferably chloro or fluoro. More preferably, R<sub>6</sub> is H or methyl.

[0054] According to another embodiment, W is -C(O)-C(O)-N(R<sub>6</sub>)<sub>2</sub>, wherein R<sub>6</sub> is hydrogen, (C1-C6)-alkyl, (C1-C6)-alkenyl, (C6-C10)-aryl-(C1-C6)-alkyl-, or (C6-C10)-heteroaryl-(C1-C6)-alkyl-, wherein R<sub>6</sub> is optionally substituted with up to 3 J groups. Preferably, R<sub>6</sub> is hydrogen, (C3-C10)-cycloalkyl or -cycloalkenyl, or (C3-C10)-heterocyclyl. Alternatively, one R<sub>6</sub> is hydrogen and the other R<sub>6</sub> is (C6-C10)-aryl-(C1-C3)alkyl-, wherein the alkyl is optionally substituted with CO<sub>2</sub>H; (C3-C6)cycloalkyl-; (C5)-heterocyclyl-(C1-C3)alkyl-; (C3-C6)alkenyl-; or each R<sub>6</sub> is (C1-C6)-alkyl-. Alternatively, each R<sub>6</sub> is (C1-C3)-alkyl-.

[0055] Most preferably, -NHR<sub>6</sub> in W is:




Preferred J substituents on the alkyl and aryl groups in this embodiment are halogen, carboxy, and heteroaryl.

More preferred substituents on the aryl groups are


- 5 halogen (preferably chloro or fluoro) and more preferred J substituents on the alkyl groups are carboxy and heteroaryl.

[0056] According to yet other preferred embodiments of formula I, W is:




wherein the  $\text{NR}^6\text{R}^6$  is  $-\text{NH}-(\text{C1-C6 aliphatic})$ ,  $-\text{NH}-(\text{C3-C6 cycloalkyl})$ ,  $-\text{NH}-\text{CH}(\text{CH}_3)-\text{aryl}$ , or  $-\text{NH}-\text{CH}(\text{CH}_3)-\text{heteroaryl}$ , wherein said aryl or said heteroaryl is optionally substituted with up to 3 halogens.

- 15 [0057] In any preferred embodiment of W, the  $\text{NR}^6\text{R}^6$  is:




[0058] In any preferred embodiment of W, the  $\text{NR}^6\text{R}^6$  is:




5

[0059] In any more preferred embodiment of W, the  $\text{NR}^6\text{R}^6$  is:




10 [0060] In any more preferred embodiment of W, the  $\text{NR}^6\text{R}^6$  is:




15 [0061] According to one embodiment,  $\text{R}^1$  is selected from:

- 27 -




[0062] According to a preferred embodiment, R<sup>1</sup> is selected from:

5




[0063] According to another embodiment, R<sup>1</sup> is:

10




[0064] According to another embodiment, R<sup>1</sup> is:




[0065] Most preferably, R<sup>1</sup> is cyclohexyl.


15 [0066] According to one embodiment, R<sup>2</sup> is:




[0067] According to another embodiment, R<sup>2</sup> is



[0068] According to any preferred embodiments,  $R^2$  is:




5 [0069] According to any more preferred embodiments,  $R^2$  is:




[0070] In any most preferred embodiments,  $R^2$  is t-  
10 butyl.

[0071] According to one embodiment,  $R^3$  is:



[0072] According to a preferred embodiment,  $R^3$  is:




[0073] More preferably,  $R^3$  is propyl (preferably, n-propyl).

20 [0074] In any preferred embodiment of this invention,  $R^{3'}$  is H.


[0075] According to another embodiment,  $R^3$  and  $R^{3'}$  together with the atom to which they are bound form the ring system:



5 [0076] According to another embodiment,  $R^{3'}$  is hydrogen and  $R^3$  is:




[0077] According to preferred embodiments,  $R^{3'}$  is hydrogen and  $R^3$  is:



10

[0078] In yet other embodiments of this invention,  $R^{3'}$  is hydrogen and  $R^3$  is:



15 [0079] According to one embodiment, one of  $R^4$  or  $R^{4'}$  is hydrogen.

[0080] According to another embodiment, one of  $R^4$  or  $R^{4'}$  (C1-C6)-alkyl.

[0081] According to a preferred embodiment,  $R^4$  and  $R^{4'}$  is hydrogen.

20

[0082] In certain embodiments, 1 or 2 carbon atoms of  $R^1$ ,  $R^2$ ,  $R^3$ , or  $R^4$ , are optionally and independently replaced with N, NH, O, or S.

[0083] Accordingly, one embodiment of this invention provides a compound wherein  $R^1$  is cyclohexyl, wherein 1 or 2 carbon atoms is optionally replaced with N, NH, O, or S and wherein each atom is optionally and independently substituted with 1, 2, or 3 J groups, wherein J is halogen, OH, OR', NH,  $N(R')_2$  (and  $R'$  is, preferably, (C1-C6)-alkyl).

[0084] In certain other embodiments no carbon atoms of  $R^1$ ,  $R^2$ ,  $R^3$ , and  $R^4$  are replaced with N, NH, O, or S. In other embodiments these groups have no J substituents.

[0085] The compounds of this invention may contain one or more asymmetric carbon atoms and thus may occur as racemates and racemic mixtures, single enantiomers, diastereomeric mixtures and individual diastereomers. All such isomeric forms of these compounds are expressly included in the present invention. Each stereogenic carbon may be of the R or S configuration.

[0086] Preferably, the compounds of this invention have the structure and stereochemistry depicted in compounds 1-76.

[0087] Any of the preferred embodiments recited above, including those embodiments in the above species, may define formula I individually or be combined to produce a preferred embodiment of this invention.

[0088] Abbreviations which are used in the schemes, preparations and the examples that follow are:

THF: tetrahydrofuran

DMF: N,N,-dimethylformamide

EtOAc: ethyl acetate

AcOH: acetic acid

HOBT: 1-hydroxybenzotriazole hydrate


- EDC: 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide  
hydrochloride
- NMM: N-methylmorpholine
- NMP: N-methylpyrrolidinone
- 5 EtOH: ethanol
- t-BuOH: tert-butanol
- Et<sub>2</sub>O: diethyl ether
- BOC: tert-butyloxycarbonyl
- BOC<sub>2</sub>O: di-tert-butyl dicarbonate
- 10 Cbz: benzyloxycarbonyl
- Chg: cyclohexylglycine
- TbG: tert-butylglycine
- Fmoc: 9-fluorenyl methyloxycarbonyl
- DMSO: dimethyl sulfoxide
- 15 TFA: trifluoroacetic acid
- DCCA: dichloroacetic acid
- DCE: dichloroethane
- DIEA: diisopropylethylamine
- MeCN: acetonitrile
- 20 PyBrOP: tris(pyrrolidino)bromophosphonium  
hexafluorophosphate
- TBTU or HATU: 2-(1H-benzotriazole-1-yl)-1,1,3,3-  
tetramethyluronium tetrafluoroborate
- DMAP: 4-dimethylaminopyridine
- 25 PPTS: pyridinium p-toluenesulfonate
- IBX: periodobenzoic acid
- AIBN: 2,2'-azobisisobutyronitrile
- TEMPO: 2,2,6,6-tetramethyl-1-piperidinyloxy, free  
radical)
- 30 rt or RT: room temperature
- ON: overnight
- ND: not determined
- MS: mass spectrometry
- LC: liquid chromatography




General Synthetic Methodology:

[0089] The compounds of this invention may be prepared in general by methods known to those skilled in the art. Schemes 1-7 below illustrate synthetic routes to the compounds of the present invention. Other equivalent schemes, which will be readily apparent to the ordinary skilled organic chemist, may alternatively be used to synthesize various portions of the molecule as illustrated by the general schemes below, and the preparative examples that follow.

Scheme 1



**12**


[0090] Scheme 1 above provides a general synthetic route for the preparation of compounds of formula I with various Ar groups, wherein T is pyrazine, R<sup>1</sup> is cyclohexyl, R<sup>2</sup> is t-butyl, and W is a -C(O)C(O)-N(H)-cyclopropyl. As would be recognized by skilled

practitioners, compounds of formula I with T, R<sup>1</sup>, and R<sup>2</sup> other than those depicted above may be prepared by varying the synthetic route.

[0091] For example, compounds of formula I wherein T  
5 is other than pyrazine may be prepared by reacting compound 5, (after removal of the Z group) with a compound of formula T-C(O)-OH under appropriate coupling conditions.


[0092] Similarly, substituting another amino acid  
10 derivative for Z-Tbg-OH in the conversion of compound 3 to compound 4 or another amino acid derivative for Z-Chg-OH in the conversion of compound 4 to compound 5 would provide compounds with varying R<sup>2</sup> and R<sup>1</sup> groups, respectively.

## Scheme 2



[0093] Scheme 2 above provides a general synthetic route for the preparation of compound 59.

## 5 Scheme 3





[0094] An alternative approach to preparing compounds of this invention is depicted in Scheme 3. In this approach, a 4-hydroxyproline derivative 13 is reacted

with a compound 14 to provide a compound 15. In Scheme 3, P<sup>1</sup> is hydrogen or an appropriate amine protecting group, P<sup>2</sup> is hydrogen or an appropriate carboxy protecting group, Ar is as defined herein, and X is an appropriate  
5 leaving group. In one embodiment set forth below, P<sup>1</sup> is t-butoxycarbonyl, P<sup>2</sup> is hydrogen, Ar is 4-chloro-2-pyridine, X is Cl, and 13 and 14 are reacted in the presence of tBuOK, DMSO, and THF.

[0095] As would be appreciated by skilled  
10 practitioners, the compound 15 then may be carried on to compounds of formula I by routine methods. One such method is depicted below in Scheme 4


Scheme 4





[0096] Scheme 5 depicts an alternative approach for preparing a compound of this invention (59). The steps used in scheme 5 could be modified by, for example, using different reagents or carrying out the reactions in a different order.


#### Scheme 6





[0097] Scheme 6 depicts an approach for preparing compound 28. In this embodiment, a compound 27 is converted to compound 28 by removal of the benzyloxy carbonyl protecting group under hydrogenolysis conditions. This scheme 6 could be modified using techniques known to skilled practitioners to arrive at compound 28.


Scheme 7



10 [0098] Scheme 7 depicts an alternative approach to prepare compounds of this invention. In scheme 7, the variables are as described herein.

[0099] Accordingly, one embodiment of this invention provides a process for preparing a compound of formula I, as defined in any of the embodiments herein, comprising  
15 the step of: reacting a compound of formula II in the

presence of a compound of formula III to provide a compound of formula IV:



5 wherein:

$R^{10}$  is an amine protecting group, a P3- residue of an HCV protease inhibitor described herein, or a P4-P3- residue of an HCV protease inhibitor as described herein, and wherein the P3 and the P4-P3 residues are optionally protected the an amino-terminal capping group;

$R^{11}$  is a carboxy protecting group or a P1 residue of an HCV protease inhibitor described herein, wherein the P1 residue is optionally protected with a carboxy terminal protecting group or with W. Ar is as define in any of the embodiments herein. X is an appropriate leaving group. As would be appreciated by skilled practitioners, an appropriate leaving group may be generated in situ.

[0100] In an alternative embodiment, the 4-hydroxy group in formula II may be converted to a leaving group. In such an embodiment, X is a nucleophilic oxygen which reacts with II to provide IV.

[0101] As used herein, P1, P3, P4 refer to the residues of an HCV protease inhibitor as defined in the art and as are well known to skilled practitioners.

[0102] The compound of formula IV may be carried on to a compound of formula I according to the methods described herein.

[0103] Although certain exemplary embodiments are depicted and described below, it will be appreciated that compounds of this invention can be prepared according to the methods described generally above using appropriate

starting materials generally available to one of ordinary skill in the art.

[0104] Another embodiment of this invention provides a composition comprising a compound of formula I or a  
5 pharmaceutically acceptable salt thereof. According to a preferred embodiment, the compound of formula I is present in an amount effective to decrease the viral load in a sample or in a patient, wherein said virus encodes a serine protease necessary for the viral life cycle, and a  
10 pharmaceutically acceptable carrier.

[0105] If pharmaceutically acceptable salts of the compounds of this invention are utilized in these compositions, those salts are preferably derived from inorganic or organic acids and bases. Included among  
15 such acid salts are the following: acetate, adipate, alginate, aspartate, benzoate, benzene sulfonate, bisulfate, butyrate, citrate, camphorate, camphor sulfonate, cyclopentane-propionate, digluconate, dodecylsulfate, ethanesulfonate, fumarate,  
20 glucoheptanoate, glycerophosphate, hemisulfate, heptanoate, hexanoate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, lactate, maleate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, oxalate, pamoate, pectinate, persulfate,  
25 3-phenyl-propionate, picrate, pivalate, propionate, succinate, tartrate, thiocyanate, tosylate and undecanoate. Base salts include ammonium salts, alkali metal salts, such as sodium and potassium salts, alkaline earth metal salts, such as calcium and magnesium salts,  
30 salts with organic bases, such as dicyclohexylamine salts, N-methyl-D-glucamine, and salts with amino acids such as arginine, lysine, and so forth.

[0106] Also, the basic nitrogen-containing groups may be quaternized with such agents as lower alkyl halides,

such as methyl, ethyl, propyl, and butyl chloride, bromides and iodides; dialkyl sulfates, such as dimethyl, diethyl, dibutyl and diamyl sulfates, long chain halides such as decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides, aralkyl halides, such as benzyl and phenethyl bromides and others. Water or oil-soluble or dispersible products are thereby obtained.

[0107] The compounds utilized in the compositions and methods of this invention may also be modified by appending appropriate functionalities to enhance selective biological properties. Such modifications are known in the art and include those which increase biological penetration into a given biological system (e.g., blood, lymphatic system, central nervous system), increase oral availability, increase solubility to allow administration by injection, alter metabolism and alter rate of excretion.

[0108] Pharmaceutically acceptable carriers that may be used in these compositions include, but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, polyethylene glycol, sodium carboxymethylcellulose, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, polyethylene glycol and wool fat.

[0109] According to a preferred embodiment, the compositions of this invention are formulated for

pharmaceutical administration to a mammal, preferably a human being.

[0110] Such pharmaceutical compositions of the present invention may be administered orally, parenterally, by  
5 inhalation spray, topically, rectally, nasally, buccally, vaginally or via an implanted reservoir. The term "parenteral" as used herein includes subcutaneous, intravenous, intramuscular, intra-articular, intra-synovial, intrasternal, intrathecal, intrahepatic,  
10 intralesional and intracranial injection or infusion techniques. Preferably, the compositions are administered orally or intravenously.

[0111] Sterile injectable forms of the compositions of this invention may be aqueous or oleaginous suspension.  
15 These suspensions may be formulated according to techniques known in the art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally  
20 acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed  
25 as a solvent or suspending medium. For this purpose, any bland fixed oil may be employed including synthetic mono- or di-glycerides. Fatty acids, such as oleic acid and its glyceride derivatives are useful in the preparation of injectables, as are natural pharmaceutically-  
30 acceptable oils, such as olive oil or castor oil, especially in their polyoxyethylated versions. These oil solutions or suspensions may also contain a long-chain alcohol diluent or dispersant, such as carboxymethyl cellulose or similar dispersing agents which are commonly

used in the formulation of pharmaceutically acceptable dosage forms including emulsions and suspensions. Other commonly used surfactants, such as Tweens, Spans and other emulsifying agents or bioavailability enhancers which are commonly used in the manufacture of pharmaceutically acceptable solid, liquid, or other dosage forms may also be used for the purposes of formulation.

[0112] Dosage levels of between about 0.01 and about 100 mg/kg body weight per day, preferably between about 0.5 and about 75 mg/kg body weight per day of the protease inhibitor compounds described herein are useful in a monotherapy for the prevention and treatment of antiviral, particularly anti-HCV mediated disease. Typically, the pharmaceutical compositions of this invention will be administered from about 1 to about 5 times per day or alternatively, as a continuous infusion. Such administration can be used as a chronic or acute therapy. The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. A typical preparation will contain from about 5% to about 95% active compound (w/w). Preferably, such preparations contain from about 20% to about 80% active compound. As recognized by skilled practitioners, dosages of interferon are typically measured in IU (e.g., about 4 million IU to about 12 million IU).

[0113] When the compositions of this invention comprise a combination of a compound of formula I and one or more additional therapeutic or prophylactic agents, both the compound and the additional agent should be present at dosage levels of between about 10 to 100%, and

more preferably between about 10 to 80% of the dosage normally administered in a monotherapy regimen.

[0114] The pharmaceutical compositions of this invention may be orally administered in any orally acceptable dosage form including, but not limited to, capsules, tablets, aqueous suspensions or solutions. In the case of tablets for oral use, carriers that are commonly used include lactose and corn starch. Lubricating agents, such as magnesium stearate, are also typically added. For oral administration in a capsule form, useful diluents include lactose and dried cornstarch. When aqueous suspensions are required for oral use, the active ingredient is combined with emulsifying and suspending agents. If desired, certain sweetening, flavoring or coloring agents may also be added.

[0115] Alternatively, the pharmaceutical compositions of this invention may be administered in the form of suppositories for rectal administration. These may be prepared by mixing the agent with a suitable non-irritating excipient which is solid at room temperature but liquid at rectal temperature and therefore will melt in the rectum to release the drug. Such materials include cocoa butter, beeswax and polyethylene glycols.

[0116] The pharmaceutical compositions of this invention may also be administered topically, especially when the target of treatment includes areas or organs readily accessible by topical application, including diseases of the eye, the skin, or the lower intestinal tract. Suitable topical formulations are readily prepared for each of these areas or organs.

[0117] Topical application for the lower intestinal tract may be effected in a rectal suppository formulation

(see above) or in a suitable enema formulation.

Topically-transdermal patches may also be used.

[0118] For topical applications, the pharmaceutical compositions may be formulated in a suitable ointment  
5 containing the active component suspended or dissolved in one or more carriers. Carriers for topical administration of the compounds of this invention include, but are not limited to, mineral oil, liquid petrolatum, white petrolatum, propylene glycol,  
10 polyoxyethylene, polyoxypropylene compound, emulsifying wax and water. Alternatively, the pharmaceutical compositions may be formulated in a suitable lotion or cream containing the active components suspended or dissolved in one or more pharmaceutically acceptable  
15 carriers. Suitable carriers include, but are not limited to, mineral oil, sorbitan monostearate, polysorbate 60, cetyl esters wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water.

[0119] For ophthalmic use, the pharmaceutical  
20 compositions may be formulated as micronized suspensions in isotonic, pH adjusted sterile saline, or, preferably, as solutions in isotonic, pH adjusted sterile saline, either with or without a preservative such as benzylalkonium chloride. Alternatively, for ophthalmic  
25 uses, the pharmaceutical compositions may be formulated in an ointment such as petrolatum.


[0120] The pharmaceutical compositions of this invention may also be administered by nasal aerosol or inhalation. Such compositions are prepared according to  
30 techniques well known in the art of pharmaceutical formulation and may be prepared as solutions in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability,



fluorocarbons, and/or other conventional solubilizing or dispersing agents.

[0121] Most preferred are pharmaceutical compositions formulated for oral administration.

- 5 [0122] In another embodiment, the compositions of this invention additionally comprise another anti-viral agent, preferably an anti-HCV agent. Such anti-viral agents include, but are not limited to, immunomodulatory agents, such as  $\alpha$ -,  $\beta$ -, and  $\gamma$ -interferons, pegylated derivatized
- 10 interferon- $\alpha$  compounds, and thymosin; other anti-viral agents, such as ribavirin, amantadine, and telbivudine; other inhibitors of hepatitis C proteases (NS2-NS3 inhibitors and NS3-NS4A inhibitors); inhibitors of other targets in the HCV life cycle, including helicase and
- 15 polymerase inhibitors; inhibitors of internal ribosome entry; broad-spectrum viral inhibitors, such as IMPDH inhibitors (e.g., compounds of United States Patent 5,807,876, 6,498,178, 6,344,465, 6,054,472, WO 97/40028, WO 98/40381, WO 00/56331, and mycophenolic acid and
- 20 derivatives thereof, and including, but not limited to VX-497, VX-148, and/or VX-944); or combinations of any of the above. See also W. Markland et al., Antimicrobial & Antiviral Chemotherapy, 44, p. 859 (2000) and U.S. Patent 6,541,496.



VX-497

[0123] The following definitions are used herein (with trademarks referring to products available as of this application's filing date).

- [0124] "Peg-Intron" means PEG-Intron®, peginterferon alfa-2b, available from Schering Corporation, Kenilworth, NJ;
- [0125] "Intron" means Intron-A®, interferon alfa-2b  
5 available from Schering Corporation, Kenilworth, NJ;
- [0126] "ribavirin" means ribavirin (1-beta-D-ribofuranosyl-1H-1,2,4-triazole-3-carboxamide, available from ICN Pharmaceuticals, Inc., Costa Mesa, CA; described in the Merck Index, entry 8365, Twelfth Edition; also  
10 available as Rebetol® from Schering Corporation, Kenilworth, NJ, or as Copegus® from Hoffmann-La Roche, Nutley, NJ;
- [0127] "Pagasys" means Pagasys®, peginterferon alfa-2a available Hoffmann-La Roche, Nutley, NJ;
- 15 [0128] "Roferon" mean Roferon®, recombinant interferon alfa-2a available from Hoffmann-La Roche, Nutley, NJ;
- [0129] "Berefor" means Berefor®, interferon alfa 2 available from Boehringer Ingelheim Pharmaceutical, Inc., Ridgefield, CT;
- 20 [0130] Sumiferon®, a purified blend of natural alpha interferons such as Sumiferon available from Sumitomo, Japan;
- [0131] Wellferon®, interferon alpha n1 available from Glaxo\_Wellcome LTd., Great Britain;
- 25 [0132] Alferon®, a mixture of natural alpha interferons made by Interferon Sciences, and available from Purdue Frederick Co., CT;
- [0133] The term "interferon" as used herein means a member of a family of highly homologous species-specific  
30 proteins that inhibit viral replication and cellular proliferation, and modulate immune response, such as interferon alpha, interferon beta, or interferon gamma. The Merck Index, entry 5015, Twelfth Edition.

[0134] According to one embodiment of the present invention, the interferon is  $\alpha$ -interferon. According to another embodiment, a therapeutic combination of the present invention utilizes natural alpha interferon 2a.

5 Or, the therapeutic combination of the present invention utilizes natural alpha interferon 2b. In another embodiment, the therapeutic combination of the present invention utilizes recombinant alpha interferon 2a or 2b. In yet another embodiment, the interferon is pegylated

10 alpha interferon 2a or 2b. Interferons suitable for the present invention include:

- (a) Intron (interferon-alpha 2B, Schering Plough),
- (b) Peg-Intron,
- (c) Pegasys,
- 15 (d) Roferon,
- (e) Berofer,
- (f) Sumiferon,
- (g) Wellferon,
- (h) consensus alpha interferon available from Amgen,
- 20 Inc., Newbury Park, CA,
- (i) Alferon;
- (j) Viraferon®;
- (k) Infergen®.

[0135] As is recognized by skilled practitioners, a

25 protease inhibitor would be preferably administered orally. Interferon is not typically administered orally. Nevertheless, nothing herein limits the methods or combinations of this invention to any specific dosage forms or regime. Thus, each component of a combination

30 according to this invention may be administered separately, together, or in any combination thereof.

[0136] In one embodiment, the protease inhibitor and interferon are administered in separate dosage forms. In one embodiment, any additional agent is administered as

part of a single dosage form with the protease inhibitor or as a separate dosage form. As this invention involves a combination of compounds, the specific amounts of each compound may be dependent on the specific amounts of each other compound in the combination. As recognized by skilled practitioners, dosages of interferon are typically measured in IU (e.g., about 4 million IU to about 12 million IU).

[0137] Accordingly, agents (whether acting as an immunomodulatory agent or otherwise) that may be used in combination with a compound of this invention include, but are not limited to, interferon-alpha 2B (Intron A, Schering Plough); Rebatron (Schering Plough, Inteferon-alpha 2B + Ribavirin); pegylated interferon alpha (Reddy, K.R. et al. "Efficacy and Safety of Pegylated (40-kd) interferon alpha-2a compared with interferon alpha-2a in noncirrhotic patients with chronic hepatitis C (Hepatology, 33, pp. 433-438 (2001); consensus interferon (Kao, J.H., et al., "Efficacy of Consensus Interferon in the Treatment of Chronic Hepatitis" J. Gastroenterol. Hepatol. 15, pp. 1418-1423 (2000), interferon-alpha 2A (Roferon A; Roche), lymphoblastoid or "natural" interferon; interferon tau (Clayette, P. et al., "IFN-tau, A New Interferon Type I with Antiretroviral activity" Pathol. Biol. (Paris) 47, pp. 553-559 (1999); interleukin 2 (Davis, G.L. et al., "Future Options for the Management of Hepatitis C." Seminars in Liver Disease, 19, pp. 103-112 (1999); Interleukin 6 (Davis et al. "Future Options for the Management of Hepatitis C." Seminars in Liver Disease 19, pp. 103-112 (1999); interleukin 12 (Davis, G.L. et al., "Future Options for the Management of Hepatitis C." Seminars in Liver Disease, 19, pp. 103-112 (1999); Ribavirin; and compounds that enhance the development of type 1 helper T cell

- response (Davis et al., "Future Options for the Management of Hepatitis C." Seminars in Liver Disease, 19, pp. 103-112 (1999). Interferons may ameliorate viral infections by exerting direct antiviral effects and/or by
- 5 modifying the immune response to infection. The antiviral effects of interferons are often mediated through inhibition of viral penetration or uncoating, synthesis of viral RNA, translation of viral proteins, and/or viral assembly and release.
- 10 [0138] Compounds that stimulate the synthesis of interferon in cells (Tazulakhova, E.B. et al., "Russian Experience in Screening, analysis, and Clinical Application of Novel Interferon Inducers" J. Interferon Cytokine Res., 21 pp. 65-73) include, but are not limited
- 15 to, double stranded RNA, alone or in combination with tobramycin, and Imiquimod (3M Pharmaceuticals; Sauder, D.N. "Immunomodulatory and Pharmacologic Properties of Imiquimod" J. Am. Acad. Dermatol., 43 pp. S6-11 (2000).
- [0139] Other non-immunomodulatory or immunomodulatory
- 20 compounds may be used in combination with a compound of this invention including, but not limited to, those specified in WO 02/18369, which is incorporated herein by reference (see, e.g., page 273, lines 9-22 and page 274, line 4 to page 276, line 11).
- 25 [0140] This invention may also involve administering a cytochrome P450 monooxygenase inhibitor. CYP inhibitors may be useful in increasing liver concentrations and/or increasing blood levels of compounds that are inhibited by CYP.
- 30 [0141] If an embodiment of this invention involves a CYP inhibitor, any CYP inhibitor that improves the pharmacokinetics of the relevant NS3/4A protease may be used in a method of this invention. These CYP inhibitors include, but are not limited to, ritonavir (WO 94/14436),

ketoconazole, troleandomycin, 4-methyl pyrazole, cyclosporin, clomethiazole, cimetidine, itraconazole, fluconazole, miconazole, fluvoxamine, fluoxetine, nefazodone, sertraline, indinavir, nelfinavir, 5 amprenavir, fosamprenavir, saquinavir, lopinavir, delavirdine, erythromycin, VX-944, and VX-497. Preferred CYP inhibitors include ritonavir, ketoconazole, troleandomycin, 4-methyl pyrazole, cyclosporin, and clomethiazole. For preferred dosage forms of ritonavir, 10 see United States Patent 6,037, 157, and the documents cited therein: United States Patent 5,484,801, United States Application 08/402,690, and International Applications WO 95/07696 and WO 95/09614).

[0142] Methods for measuring the ability of a compound 15 to inhibit cytochrome P50 monooxygenase activity are known (see US 6,037,157 and Yun, et al. Drug Metabolism & Disposition, vol. 21, pp. 403-407 (1993)).

[0143] Upon improvement of a patient's condition, a maintenance dose of a compound, composition or 20 combination of this invention may be administered, if necessary. Subsequently, the dosage or frequency of administration, or both, may be reduced, as a function of the symptoms, to a level at which the improved condition is retained when the symptoms have been alleviated to the 25 desired level, treatment should cease. Patients may, however, require intermittent treatment on a long-term basis upon any recurrence of disease symptoms.

[0144] It should also be understood that a specific dosage and treatment regimen for any particular patient 30 will depend upon a variety of factors, including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, rate of excretion, drug combination, and the judgment of the treating physician and the severity

of the particular disease being treated. The amount of active ingredients will also depend upon the particular described compound and the presence or absence and the nature of the additional anti-viral agent in the composition.

[0145] According to another embodiment, the invention provides a method for treating a patient infected with a virus characterized by a virally encoded serine protease that is necessary for the life cycle of the virus by administering to said patient a pharmaceutically acceptable composition of this invention. Preferably, the methods of this invention are used to treat a patient suffering from a HCV infection. Such treatment may completely eradicate the viral infection or reduce the severity thereof. More preferably, the patient is a human being.

[0146] In an alternate embodiment, the methods of this invention additionally comprise the step of administering to said patient an anti-viral agent preferably an anti-HCV agent. Such anti-viral agents include, but are not limited to, immunomodulatory agents, such as  $\alpha$ -,  $\beta$ -, and  $\gamma$ -interferons, pegylated derivatized interferon- $\alpha$  compounds, and thymosin; other anti-viral agents, such as ribavirin and amantadine; other inhibitors of hepatitis C proteases (NS2-NS3 inhibitors and NS3-NS4A inhibitors); inhibitors of other targets in the HCV life cycle, including helicase and polymerase inhibitors; inhibitors of internal ribosome entry; broad-spectrum viral inhibitors, such as IMPDH inhibitors (the IMPDH inhibitors disclosed in United States Patent 5,807,876, mycophenolic acid and derivatives thereof); or combinations of any of the above.

[0147] Such additional agent may be administered to said patient as part of a single dosage form comprising

both a compound of this invention and an additional anti-viral agent. Alternatively the additional agent may be administered separately from the compound of this invention, as part of a multiple dosage form, wherein  
5 said additional agent is administered prior to, together with or following a composition comprising a compound of this invention.

[0148] In yet another embodiment the present invention provides a method of pre-treating a biological substance  
10 intended for administration to a patient comprising the step of contacting said biological substance with a pharmaceutically acceptable composition comprising a compound of this invention. Such biological substances include, but are not limited to, blood and components  
15 thereof such as plasma, platelets, subpopulations of blood cells and the like; organs such as kidney, liver, heart, lung, etc; sperm and ova; bone marrow and components thereof, and other fluids to be infused into a patient such as saline, dextrose, etc.

20 [0149] According to another embodiment the invention provides methods of treating materials that may potentially come into contact with a virus characterized by a virally encoded serine protease necessary for its life cycle. This method comprises the step of contacting  
25 said material with a compound according to the invention. Such materials include, but are not limited to, surgical instruments and garments (e.g. clothes, gloves, aprons, gowns, masks, eyeglasses, footwear, etc.); laboratory instruments and garments (e.g. clothes, gloves, aprons,  
30 gowns, masks, eyeglasses, footwear, etc.); blood collection apparatuses and materials; and invasive devices, such as shunts, stents, etc.

[0150] In another embodiment, the compounds of this invention may be used as laboratory tools to aid in the



isolation of a virally encoded serine protease. This method comprises the steps of providing a compound of this invention attached to a solid support; contacting said solid support with a sample containing a viral  
5 serine protease under conditions that cause said protease to bind to said solid support; and eluting said serine protease from said solid support. Preferably, the viral serine protease isolated by this method is HCV NS3-NS4A protease.

10 [0151] In order that this invention be more fully understood, the following preparative and testing examples are set forth. These examples are for the purpose of illustration only and are not to be construed as limiting the scope of the invention in any way.

15 EXAMPLES

[0152] <sup>1</sup>H-NMR spectra were recorded at 500 MHz using a Bruker AMX 500 instrument. Mass spec. samples were analyzed on a MicroMass ZQ or Quattro II mass spectrometer operated in single MS mode with electrospray  
20 ionization. Samples were introduced into the mass spectrometer using flow injection (FIA) or chromatography. Mobile phase for all mass spec. analysis consisted of acetonitrile-water mixtures with 0.2% formic acid as a modifier.

25 [0153] As used herein, the term "R<sub>t</sub>(min)" or "RT" refers to the HPLC retention time, in minutes, associated with the compound. The HPLC retention times listed were obtained from the mass spec. data or using the following method (Method B):

30 Instrument: Hewlett Packard HP-1050;  
Column: YMC C<sub>18</sub> (Cat. No. 326289C46);  
Gradient/Gradient Time: 10-90% CH<sub>3</sub>CN/H<sub>2</sub>O over 9 minutes,  
then 100% CH<sub>3</sub>CN for 2 minutes;  
Flow Rate: 0.8ml/min;

Detector Wavelength: 215nm and 245nm.

[0154] Chemical naming for selected compounds herein was accomplished using the naming program provided by CambridgeSoft Corporations ChemDraw Ultra®, version 7.0.1.

### Example 1

#### Preparation of compound 59

##### 4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-benzyl ester 2-tert-butyl ester (3)

[0155] Commercially available (Bachem) Z-Hydroxy-proline(1) (10g, 42.51mmols) was dissolved 90mls of THF(tetrahydrofuran) and was cooled to 0 °C with an ice water bath. To this was added previously prepared tert-butyl N,N'-diisopropyl-imidocarbamate(2) (27ml, 135mmol) via a dropping funnel over 30 minutes. After addition the cooling bath was removed and the reaction stirred at ambient temperature for 24 hours. The volume of the reaction was reduced and then diethyl ether was added prior to washing with saturated sodium bicarbonate, then 0.5M hydrochloric acid, then water, and finally with brine. The organic layer was dried with sodium sulfate and concentrated *in vacuo* to give 15g of crude material. Material was run through a plug of SiO<sub>2</sub> and eluted with 45% EtOAc-Hexanes to give 4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-benzyl ester 2-tert-butyl ester 3 as a colorless oil 11.0g (81%)

<sup>1</sup>H NMR (CDCl<sub>3</sub>, ppm) δ 7.35 (m, 5H), 5.2 (m, 2H), 4.3 (m, 2H), 4.65 (m, 3H), 2.35 (m, 1H), 2.1 (t, 1H), 1.35, 1.55 (rotomers, 1.45, 9H).

[0156] 1-(2-{2-Cyclohexyl-2-[(pyrazine-2-carbonyl)-amino]-acetyl-amino}-3,3-dimethyl-butyl)-4-hydroxy-pyrrolidine-2-carboxylic acid tert-butyl ester (6)

[0156] Mixed (3) in EtOH, and added catalytic amount of 10% Pd on carbon, then stirred under 1 atmosphere of

hydrogen using a balloon. After 12 hours the reaction was shown to be complete by tlc, and the catalyst was filtered and washed with EtOH. The filtrate was concentrated and dried under high vacuum to give the amine as a yellow solid, which was carried on into the next step. Z-Tbg-OH (8.3g, 31.1mmols) was dissolved in NMP and to it was added EDC (6.0g, 31.1mmols), HOBT (4.2g, 31.1mmols), DMAP (340mgs, 2.8mmols), and cooled to 0°C using and ice-water bath. To this mixture was added the amine as a solution in NMP, and the reaction was stirred for 2 days. The reaction was poured over ice and acidified with 0.5N hydrochloric acid to pH 5, and then extracted with EtOAc. The organic extracts were washed with saturated sodium bicarbonate, then water, and finally with brine. The organic layer was dried with sodium sulfate and concentrated in vacuo to give 14.8g of crude material. Purification was carried out using chromatography on SiO<sub>2</sub>, eluting with 50% EtOAc - Hexanes. Concentration of the homogeneous fractions yielded 10.5 grams of 4 as a colorless foam (85%) and used as is in the next step.

[0157] To a mixture of 4 (10.5g, 24.16mmol) in EtOH, was added a catalytic amount of 10% Pd on carbon, then stirred under 1 atmosphere of hydrogen using a balloon. After 12 hours the reaction was shown to be complete by tlc, and the catalyst was filtered and washed with EtOH. The filtrate was concentrated and dried under high vacuum to give the amine as a yellow solid, which was carried on into the next step. Z-Chg-OH (7.7g, 26.6mmols) was dissolved in NMP and to it was added EDC (5.1g, 26.7mmols), HOBT (3.6g, 26.6mmols), and cooled to 0°C using and ice-water bath. To this mixture was added the previously prepared amine as a solution in NMP, and the reaction was stirred for 2 days. The reaction was poured

over ice and brine, and then extracted with EtOAc. The organic extracts were washed with 0.5N hydrochloric acid, saturated sodium bicarbonate, water, and finally with brine. The organic extract was dried with sodium sulfate and concentrated in vacuo to give 15.31g of 5 as crude material, which was used as is in the next step.

[0158] To a solution of 5 (5.6g, 9.76mmol) in EtOH, was added a catalytic amount of 10% Pd on carbon, then stirred under 1 atmosphere of hydrogen using a balloon. After 12 hours the reaction was shown to be complete by tlc, and the catalyst was filtered and washed with EtOH. The filtrate was concentrated and dried under high vacuum to give the amine as an amorphous solid, which was carried on into the next step. Pyrazine-2-carboxylic acid (1.45g, 11.7mmols) was dissolved in NMP and to it was added EDC (2.24g, 11.7mmols), HOBT (1.34g, 11.7mmols), and cooled to 0°C using an ice bath. To this mixture was added the previously prepared amine as a solution in NMP, and the reaction was stirred for 2 days. The reaction was poured over ice and brine, and then extracted with EtOAc. The organic extracts were washed with 0.5N hydrochloric acid, saturated sodium bicarbonate, water, and finally with brine. The organic extract was dried with sodium sulfate and concentrated in vacuo to give 5.3g (99%) of 6 as a colorless foam, which was used as is in the next step.

Pyrazine-2-carboxylic acid ({1-[4-(5-chloro-pyridin-2-yloxy)-2-(1-cyclopropylaminooxalyl-butylcarbamoyl)-pyrrolidine-1-carbonyl]-2,2-dimethyl-propylcarbamoyl}-cyclohexyl-methyl)-amide (59)

To a solution of 6 (0.15g, 0.28mmols) in anhydrous THF was added triphenylphosphine (0.131g, 0.5mmols), 2-hydroxy-4-chloro-pyridine (65mg, 0.5mmols), and last was added the diethyl azodicarboxylate (0.100mL, 1.85mmols).

The reaction was stirred at room temperature for 18 hours or until the reaction showed no 6 remaining by HPLC. THF was removed from the reaction and then the material was taken up in EtOAc, and washed with 0.1N NaOH, 0.5N hydrochloric acid, water, and finally with brine. The organic extract was dried with sodium sulfate and concentrated *in vacuo* to give the crude tert-butyl ester. The tert-butyl ester group was hydrolyzed to the carboxylic acid by treatment with 50% trifluoroacetic acid in dichloromethane for 3 hours. The solvent was removed under vacuum, and then the residue was taken up with 0.1N NaOH, and washed with EtOAc. The aqueous phase was acidified with 5% citric acid, and then extracted with EtOAc. The resultant organic phase was washed with water and then brine, and then the organic extract was dried with sodium sulfate and concentrated *in vacuo* to give 4-(5-Chloro-pyridin-2-yloxy)-1-(2-(2-cyclohexyl-2-[(pyrazine-2-carbonyl)-amino]-acetylamino)-3,3-dimethylbutyryl)-pyrrolidine-2-carboxylic acid **7a** as a colorless foam, which was used as is in the next step.

To a solution of **7a** in 2mls of dimethylformamide was added TBTU (0.15g, 0.47mmols), DIEA (0.15mL, 1.1mmols), and the reaction was stirred for 1.5 hours, and then the amine **10** [U. Schoellkogf et al., Justus Liebigs Ann. Chem. GE, pp. 183-202 (1976) and J. Semple et al., Org. Letts., 2, pp. 2769-2772 (2000) was added to the mixture followed by 4-methylmorpholine (0.2mL, 1.82mmol). The reaction was stirred at ambient temperature for 12 hours, and then was poured over water and extracted EtOAc. The organic extract was dried with sodium sulfate and concentrated *in vacuo* to give pyrazine-2-carboxylic acid [(1-{4-(5-chloro-pyridin-2-yloxy)-2-[1-(cyclopropylcarbamoyl-hydroxy-methyl)-butylcarbamoyl]-pyrrolidine-1-carbonyl}-2,2-dimethyl-propylcarbamoyl)-

cyclohexyl-methyl]-amide 11a (40mgs) as a colorless foam, which was used as is in the next step. To a solution of 11a (40mg) in dichloromethane (4ml) was added tert-butanol 0.1mL, and Dess-Martin periodinane (40mgs, 5 0.086mmol), and then stirred at ambient temperature for 6 hours. To the reaction was added 1mL of a 1:1 mixture of 1N sodium thiosulfate and saturated sodium bicarbonate. After 15 minutes the reaction was extracted with EtOAc and then the solvent was removed under vacuum.

10 Purification was carried out using chromatography on SiO<sub>2</sub>, eluting with 50% EtOAc - Hexanes. Concentration of the homogeneous fractions yielded 0.095 grams of 59 as a colorless foam (4.5% based on 0.28mmols of 6). <sup>1</sup>H NMR (CDCl<sub>3</sub>, ppm) δ 9.39(s,1H), 8.76(s,1H), 8.56(s,1H), 8.24(d,1H,J=9.6Hz), 8.08(s,1H), 7.8(d,1H,6.4Hz), 7.69(s,1H), 7.47-7.40(m,2H), 7.47(d,1H,J=8.75Hz), 5.63(s,1H), 5.60-5.50(m,1H), 4.90(m,1H), 4.70(m,1H), 4.11(d,1H,J=11.6Hz), 4.0(m,1H), 2.90(m,1H) 2.60(m,1H), 2.25(m,1H), 2.0(m,1.90-1.4(m,1H), 1.25-0.8(m,18H), 20 (0.73(m,2H); LC/MS : RT = 3.61min, 4.15min (10-90% CH<sub>3</sub>CN/7min); MH<sup>+</sup> = 767.3, M<sup>-</sup> = 765.5.

#### Example 2

##### Alternative Preparation of Compound 59

[0159] Boc-Pro(4(R)- 5-Chloro-pyridin-2-yloxy)-OH

25 (15a) Boc-Hyp-OH (130 g, 562.16 mmol) was dissolved in anhydrous DMSO (1.6 L) and to this solution was added 1M potassium tert- butoxide in THF (1.4 L, 140 mmol) maintaining the internal temperature under 25 °C. After stirring the solution for 1.5 h at RT, 2,5-Dichloro-30 pyridine (90.0g, 608.15 mmol) was added and the reaction mixture was stirred for 18 h at RT. The mixture was poured into water (2.5 L) and extracted with Ethyl Ether (1 L) to remove excess 2,5-Dichloro-pyridine. The aqueous layer was then acidified with 1N HCL (0.8 L) and

extracted two times with Ethyl Acetate (2.5 L). The organic layers were combined and washed with Brine. The Ethyl Acetate was dried with magnesium sulfate, filtered and concentrated under reduced pressure to give a brown oil - 210 g crude material.

Boc-Pro (4(R)- 5-Chloro-pyridin-2-yloxy)-allyl ester (16)

[0160] Boc-Pro (4(R)- 5-Chloro-pyridin-2-yloxy)-OH (15a) (~210g, 557mmol) was dissolved in anhydrous acetonitrile (1.5 L) 2,3,4,6,7,8,9,10-Octahydro-pyrimido [1,2-a] azepine (DBU) (0.13 L, 869.28 mmol) and allyl bromide (81.5g, 673.66 mmol) were added successively and the reaction mixture was stirred for 18 h at RT. The mixture was concentrated, the resulting oil was diluted with Ethyl Acetate (2 L) and washed successively with water two times 500 mL and Brine 500 mL. The Ethyl Acetate layer was dried with magnesium sulfate, filtered and concentrated under reduced pressure to give a brown oil which was applied to a silica column with methylene chloride and eluted with 25% ethyl acetate in hexane to yield a yellow oil, 181 g, 472.77 mmol, 84% for two steps.

Pro (4(R)- 5-Chloro-pyridin-2-yloxy)-allyl ester (17)

[0161] Boc-Pro (4(R)- 5-Chloro-pyridin-2-yloxy)-allyl ester (16) (181g, 427.77 mmol) was treated with cold 0 °C (440 mL) 90:10 Trifluoroacetic acid, Methylene Chloride. The mixture was allowed to come to RT and stirred for 3 h. After 3 h 400 mL Toluene was added to the mixture and it was concentrated under reduced pressure to yield the crude Trifluoroacetic acid salt of (17).

Boc-Tbg-Pro (4(R)- 5-Chloro-pyridin-2-yloxy)-allyl ester (18)

[0162] Pro (4(R)- 5-Chloro-pyridin-2-yloxy)-allyl ester (17) (187g, theor.), the crude trifluoroacetic

acid salt was coupled to a cooled 0 °C solution of Boc-Tbg-OH (110g, 475.56 mmol), NMM (155 mL, 1,410 mmol), EDC (99g, 518.32 mmol), HOBt (70 g, 518.32 mmol) in 400 mL Methylene Chloride. The mixture was allowed to come to  
5 RT and stirred for 18 h. The mixture was concentrated, the resulting oil was diluted with Ethyl Acetate (2 L) and washed successively with 0.5N HCL two times 500 mL, and water 500 mL, Brine 500 mL. The Ethyl Acetate layer was dried with magnesium sulfate, filtered and  
10 concentrated under reduced pressure to give a brown oil which was applied to a silica column with methylene chloride and eluted with 15% ethyl acetate in hexane to yield a pale yellow foam, 180 g, 362.83 mmol, 77%.  
Tbg-Pro (4(R)- 5-Chloro-pyridin-2-yloxy)-allyl ester  
15 (19)

[0163] Boc-Tbg-Pro (4(R)- 5-Chloro-pyridin-2-yloxy)-allyl ester (18) (180g, 362.83 mmol), was treated with cold 0 °C (330 mL) 90:10 Trifluoroacetic acid, Methylene Chloride. The mixture was allowed to come to RT and  
20 stirred for 3 h. After 3 h Toluene (200 mL) was added to the mixture and it was concentrated under reduced pressure to yield a yellow oil to which methylene chloride (100 mL), ethyl ether (1.5 L) were added successively. The mixture was stirred and 4N HCL dioxane  
25 was added (50 mL) stirring was continued for 1h and the crude HCL dipeptide salt was filtered and washed with cold ethyl ether 0 °C to yield 142.5g, 323.86 mmol 91%.  
Boc-Chg-Tbg-Pro (4(R)- 5-Chloro-pyridin-2-yloxy)-allyl ester (20)

30 [0164] Tbg-Pro (4(R)- 5-Chloro-pyridin-2-yloxy)-allyl ester (19) (142.5g, 323.86 mmol) was coupled to a cooled 0 °C solution of Boc-Chg-OH (94g, 365.29 mmol), NMM (109 mL, 994.18 mmol), EDC (69g, 361.26 mmol), HOBt (48.77 g, 361.26 mmol) in 400 mL Methylene Chloride. The mixture



was allowed to come to RT and stirred for 18 h. The mixture was concentrated, the resulting oil was diluted with Ethyl Acetate (2 L) and washed successively with 0.5N HCL two times 500 mL, and water 500 mL, Brine 500 mL. The Ethyl Acetate layer was dried with magnesium sulfate, filtered and concentrated under reduced pressure to give a brown oil which was applied to a silica column with methylene chloride and eluted with 25% ethyl acetate in hexane to yield a pale yellow foam, 180 g, 362.83 mmol, 77%.

Chg-Tbg-Pro (4(R)- 5-Chloro-pyridin-2-yloxy)-allyl ester (21)

[0165] Boc-Chg-Tbg-Pro (4(R)- 5-Chloro-pyridin-2-yloxy)-allyl ester (20) (65 g, 102.33 mmol) was treated with cold 0 °C (250 mL) 90:10 Trifluoroacetic acid, Methylene Chloride. The mixture was allowed to come to RT and stirred for 3 h. After 3 h Toluene (200 mL) was added to the mixture and it was concentrated under reduced pressure to yield a yellow oil to which methylene chloride (200 mL) was added the mixture was stirred and, ethyl ether (1.5 L) were added successively and the crude TFA tripeptide salt was filtered and washed with cold ethyl ether 0 °C to yield 66g, 101.53 mmol 98%.

Pyrazine-2- carbonyl -Chg-Tbg-Pro (4(R)- 5-Chloro-pyridin-2-yloxy)-allyl ester (22)

[0166] Chg-Tbg-Pro (4(R)- 5-Chloro-pyridin-2-yloxy)-allyl ester (21) (66g, 101.53 mmol), the crude TFA salt was coupled to a cooled 0 °C solution of Pyrazine-2-carboxylic acid (13.6g, 109.69 mmol), NMM (44 mL, 400.19 mmol), EDC (21g, 109.95 mmol), HOBt (14.85 g, 109.95 mmol) in 500 mL Methylene Chloride. The mixture was allowed to come to RT and stirred for 18 h. The mixture was concentrated, the resulting oil was diluted with Ethyl Acetate (2 L) and washed successively with 0.5N HCL

two times 500 mL, and water 500 mL, Brine 500 mL. The Ethyl Acetate layer was dried with magnesium sulfate, filtered and concentrated under reduced pressure to give a brown oil which was applied to a silica column with  
5 methylene chloride and eluted with 45% ethyl acetate in hexane to yield a pale yellow foam, 64.66 g, 100.85 mmol, 99%.

Pyrazine-2- carbonyl -Chg-Tbg-Pro (4(R)- 5-Chloro-pyridin-2-yloxy)-OH (23)

10 [0167] Pyrazine-2- carbonyl -Chg-Tbg-Pro (4(R)- 5-Chloro-pyridin-2-yloxy)-allyl ester (22) (64.66g, 100.85 mmol) was dissolved in an anhydrous mixture (250 mL) 50:50 Acetonitrile, Methylene Chloride. Tetrakis (triphenylphosphine)-palladium (0) catalyst (1.5g, 1.30  
15 mmol) was added followed by pyrrolidine (8.55 mL, 102.44 mmol). The reaction mixture was stirred at RT for 18 h. After 18 h the solvent was evaporated. The oil was dissolved in Ethyl Acetate (2L) and extracted with 10% citric acid (250 mL) two times, Brine (250 mL). The  
20 Ethyl Acetate layer was dried with magnesium sulfate, filtered and concentrated under reduced pressure to give a pale yellow solid 58.5g, 97.32 mmol 96%.

3-Amino-2-hydroxy-hexanoic acid cyclopropylamide (24)

[0168] [1-(Cyclopropylcarbamoyl-hydroxy-methyl)-  
25 butyl]-carbamic acid benzyl ester (48g, 149.82 mmol) was dissolved in Methanol (1L) and degassed with nitrogen for five minutes, palladium, 10 wt. % on activated carbon (2.5 g) was added, hydrogen was then added for 18 h. After 18 h the hydrogen was removed and the reaction was  
30 degassed with nitrogen and filtered the resulting filtrate was evaporated and dried under high vacuum to give a white solid 26.9 g, 144.42 mmol 97%.

Pyrazine-2- carbonyl -Chg-Tbg-Pro (4(R)- 5-Chloro-pyridin-2-yloxy)-Nva-hydroxy cyclopropylamide (25)

[0169] 3-Amino-2-hydroxy-hexanoic acid cyclopropylamide (24) 20g, 107.37 mmol, was coupled to Pyrazine-2-carbonyl-Chg-Tbg-Pro (4(R)-5-Chloro-pyridin-2-yloxy)-OH (23) 58.5g, 97.32 mmol, NMM (11.76 mL, 106.96 mmol), EDC (20.45g, 107.07 mmol), HOBt (14.45 g, 107.07 mmol) in 250 mL Methylene Chloride. The mixture was stirred for 18 h. The mixture was concentrated, the resulting oil was diluted with Ethyl Acetate (2 L) and washed successively with 0.5N HCL two times 500 mL, and water 500 mL, Brine 500 mL. The Ethyl Acetate layer was dried with magnesium sulfate, filtered and concentrated under reduced pressure to give a brown oil which was applied to a silica column with methylene chloride and eluted with 2% Methanol in Ethyl Acetate to yield a pale yellow foam, 61.5 g, 79.94 mmol, 74%.

Pyrazine-2-carboxylic acid ({1-[4-(5-chloro-pyridin-2-yloxy)-2-(1-cyclopropylaminooxalyl)-butylcarbamoyl]-pyrrolidine-1-carbonyl}-2,2-dimethyl-propylcarbamoyl)-cyclohexyl-methyl)-amide (59)

[0170] To a 4L round bottom equipped with an overhead stirrer a thermocouple and an active nitrogen inlet was added EDC (229.0g, 151.83 mmol), followed by anhydrous Ethyl Acetate 1230 mL, Stirring was started to effect a thick slurry. To this was added Pyrazine-2-carbonyl-Chg-Tbg-Pro (4(R)-5-Chloro-pyridin-2-yloxy)-Nva-hydroxy cyclopropylamide (25) 61.5 g, 79.94 mmol, dissolved Ethyl Acetate (250 mL) then anhydrous DMSO (460 mL) was added. A cooling bath was applied to bring the internal temperature to 7 °C. A cooled 7 °C solution of Dichloroacetic acid (65.94 mL, 787.20 mmol) in Ethyl Acetate (100 mL) was added such that the internal temperature was maintained between 12 and 25 °C. The cooling bath removed and the thin slurry was stirred for 1 h. A cooling bath was applied and the reaction was

quenched with 1N HCL (1,230 mL) while maintaining the temperature between 15 and 25 °C. The organic layer was separated and washed with water (200 mL) three times, brine (200 mL). The Ethyl Acetate layer was dried with  
5 magnesium sulfate, filtered and concentrated under reduced pressure to with a bath temperature of not more than 40 °C to give a brown oil which was applied to a silica column with methylene chloride and eluted with 90% Ethyl Acetate in hexane to yield a pale yellow foam, 44.0  
10 g, 57.34 mmol, 72%. <sup>1</sup>H NMR (CDCl<sub>3</sub>) 9.39 1H(s), 8.76 1H(s), 8.55 1H(s), 8.22 1H(d), 8.08 1H(s), 7.47 1H(NH), 7.45 1H(d) 7.35 1H(NH), 7.01 1H(NH), 6.49 1H(d), 5.63 1H(m), 5.53 1H(m), 4.88 1H(m), 4.70 1H(d), 4.67 1H(m), 4.11 1H(m), 4.01 1H(m), 2.86 1H(m), 2.57 1H(m), 2.25 1H(m),  
15 1.96 1H(m), 1.80 1H(m), 1.70 6H(m), 1.60 2H(m), 1.50 2H(m), 1.25 3H(m), 0.96 12H(m), 0.91 2H(m), 0.72 2H(m).

### Example 3

[0171] Compounds 1-72 and 74-76 have been prepared substantially as described herein. Analytical data for  
20 these compounds were consistent with the disclosed structures of the compounds. Further selected data is provided below:

LC Mass Spectrometry Data

LC Mass Spectrometry Data

Cmpd no.	Method	MASS +	RT
1	A	801.81	1.54
2	A	800.83	1.68
3	A	800.51	1.77
4	A	750.54	1.63
5	A	800.31	1.50
6	A	800.43	1.61
7	A	768.27	1.93
8	A	768.31	1.91

Cmpd no.	Method	MASS +	RT
9	A	816.26	2.07
10	A	768.28	1.99
11	A	868.27	2.18
12	A	800.26	2.16
13	A	800.17	2.03
14	A	816.14	2.07
15	A	808.16	2.17
16	A	816.12	2.25
17	A	808.41	2.09
18	A	750.39	1.97
19	A	808.40	2.17
20	A	766.34	1.99
21	A	762.43	1.94
22	A	800.38	2.08
23	A	792.30	1.87
24	A	792.30	1.97
25	A	767.67	1.43
26	A	778.00	1.48
27	A	(M+Na <sup>+</sup> ) / 824.58	1.48
28	A	(M+Na <sup>+</sup> ) / 832.65	1.46
29	A	(M+Na <sup>+</sup> ) / 824.61	1.52
30	A	(M+Na <sup>+</sup> ) / 824.61	1.51
31	A	(M+Na <sup>+</sup> ) / 856.67	1.58
32	A	(M+Na <sup>+</sup> ) / 774.63	1.39
33	A	(M+Na <sup>+</sup> ) / 800.62	1.48
34	B	734.10	3.73
52	B	790.30	3.86
53	B	800.10	4.48
54	B	768.20	4.13
55	B	768.90	4.51
56	B	766.80	4.61
57	B	778.40	3.98




Cmpd no.	Method	MASS +	RT
58	B	733.40	3.71
59	B	767.30	4.02
60	B	852.90	4.70
61	B	818.40	4.21
62	B	767.22	1.68
63	B	767.29	4.21
66	B	766.30	4.26
67	B	851.30	3.43
68	B	768.30	4.10
69	B	789.90	3.94
70	B	978.00	4.14
71	B	844.40	2.50
72	B	797.14	4.77
74	A	802.62	1.52
75	B	768.30	5.20
76	B	766.30	5.02

RT - retention time.




Method A: Hypersil BDS C18 column 5um, 2.1x50mm Flow  
rate: 1.0 ml/min Run time: 2.39 min Solvents: 0-95% MeCN.




Method B: see above.








No.	Structure
4	
5	
6	








No.	Structure
7	 <chem>CC[C@H](NC(=O)N1CC[C@H](C1)C2=CC=C(C=C2)OC3=CC=C(C=C3)F)C(=O)N[C@@H](C(C)(C)C)C(=O)N[C@@H](C4CCCCC4)C(=O)N5C=CC=CC=C5</chem>
8	 <chem>CC[C@H](NC(=O)N1CC[C@H](C1)C2=CC=C(C=C2)OC3=CC(=CC=C3)F)C(=O)N[C@@H](C(C)(C)C)C(=O)N[C@@H](C4CCCCC4)C(=O)N5C=CC=CC=C5</chem>
9	 <chem>CC[C@H](NC(=O)N1CC[C@H](C1)C2=CC=C(C=C2)OC3=CC=C(C=C3)OC(F)F)C(=O)N[C@@H](C(C)(C)C)C(=O)N[C@@H](C4CCCCC4)C(=O)N5C=CC=CC=C5</chem>




No.	Structure
10	 <chem>CC[C@H](NC(=O)C1CCN(C1)C(=O)N[C@@H](C(C)(C)C)C(=O)N[C@@H](C2CCCCC2)C(=O)Nc3ccncc3)C(=O)N[C@@H](C)C(=O)Nc4cc(F)c(F)cc4O</chem>
11	 <chem>CC[C@H](NC(=O)C1CCN(C1)C(=O)N[C@@H](C(C)(C)C)C(=O)N[C@@H](C2CCCCC2)C(=O)Nc3cc(F)(F)cc(F)c3O)C(=O)N[C@@H](C)C(=O)Nc4cc(F)(F)cc(F)c4</chem>
12	 <chem>CC[C@H](NC(=O)C1CCN(C1)C(=O)N[C@@H](C(C)(C)C)C(=O)N[C@@H](C2CCCCC2)C(=O)Nc3cc(Cl)cc(Cl)c3O)C(=O)N[C@@H](C)C(=O)Nc4ccncc4</chem>



No.	Structure
13	
14	

No.	Structure
15	
16	
17	




No.	Structure
18	
19	
20	

No.	Structure
21	
22	

No.	Structure
23	
24	
25	

No.	Structure
26	 <chem>CC(C)C(=O)N[C@@H](C(C)(C)C)C(=O)N[C@@H](C1CCCCC1)C(=O)Nc2ccncc2C3=CC=CC=C3[N+](=O)[O-]</chem>
27	 <chem>CC(C)C(=O)N[C@@H](C(C)(C)C)C(=O)N[C@@H](C1CCCCC1)C(=O)Nc2ccncc2C3=CC=CC=C3C(F)(F)F</chem>









No.	Structure
28	
29	
30	




No.	Structure
31	
32	
33	



No.	Structure
37	 <chem>CC1(C)C(=O)N(C1C(=O)N[C@@H](C2=CC=CC=C2)OCC3=CC=CC=C3)[C@H](C2=CC=CC=C2)C(=O)N[C@@H](C4=CC=CC=CC4)C(=O)Nc5cc[nH]5</chem>
38	 <chem>CC1(C)C(=O)N(C1C(=O)N[C@@H](C2=CC=CC=C2)OCC3=CC=CC=C3)C(=O)N[C@@H](C4=CC=CC=CC4)C(=O)Nc5cc[nH]5</chem>



No.	Structure
39	
40	
41	




No.	Structure
42	 <chem>CC[C@H](NC(=O)N[C@@H](C1CCCCC1)C(=O)N[C@@H](C(C)(C)C)C(=O)N2CCCC2OC3=NC4=CC=CC=C4N=CN=C35C=CC=CC=C5)C(=O)N[C@@H](C)C(=O)N6CC6</chem>
43	 <chem>CC[C@H](NC(=O)N[C@@H](C1CCCCC1)C(=O)N[C@@H](C(C)(C)C)C(=O)N2CCCC2OC3=CC=C(C=C3)F)C(=O)N[C@@H](C)C(=O)N4CC4</chem>
44	 <chem>CC[C@H](NC(=O)N[C@@H](C1CCCCC1)C(=O)N[C@@H](C(C)(C)C)C(=O)N2CCCC2OC3=CC=C(C=C3)[N+](=O)[O-])C(=O)N[C@@H](C)C(=O)N4CC4</chem>




No.	Structure
45	
46	
47	













No.	Structure
50	
51	




No.	Structure
52	
53	
54	



No.	Structure
55	
56	
57	

No.	Structure
58	
59	
60	



No.	Structure
61	
62	



No.	Structure
63	
64	
65	

No.	Structure
66	 <chem>CC(C)(C)[C@H](NC(=O)N[C@@H](C1CCCCC1)C(=O)Nc2ccncc2)[C@@H](C3CCN(C3)C(=O)N[C@@H](C[C@H](C)C(=O)Nc4ccncc4)C(=O)Oc5ccc(Cl)cc5)C(=O)Nc6ccncc6</chem>
67	 <chem>CC(C)(C)[C@H](NC(=O)N[C@@H](C1CCCCC1)C(=O)Nc2ccncc2)[C@@H](C3CCN(C3)C(=O)N[C@@H](C[C@H](C)C(=O)Nc4ccncc4)C(=O)Oc5cnc6cc(C(F)(F)F)ccc65)C(=O)Nc7ccncc7</chem>
68	 <chem>CC(C)(C)[C@H](NC(=O)N[C@@H](C1CCCCC1)C(=O)Nc2ccncc2)[C@@H](C3CCN(C3)C(=O)N[C@@H](C[C@H](C)C(=O)Nc4ccncc4)C(=O)Oc5cc(F)c(F)cc5)C(=O)Nc6ccncc6</chem>

No.	Structure
69	
70	



No.	Structure
71	
72	

No.	Structure
73	 <chem>CC1(C)C(=O)N[C@@H](C1)C(=O)N[C@@H](C2CCCCC2)C(=O)Nc3ccncc3C(=O)N[C@@H](CC)C(=O)N[C@@H](C4CC[C@H]4)C(=O)N[C@@H](C5CC[C@H]5)C(=O)Nc6cc7cc(OC)ccc7n6c8ccccc8</chem>
74	 <chem>CC1(C)C(=O)N[C@@H](C1)C(=O)N[C@@H](C2CCCCC2)C(=O)Nc3ccncc3C(=O)N[C@@H](CC)C(=O)N[C@@H](C4CC[C@H]4)C(=O)N[C@@H](C5CC[C@H]5)C(=O)Nc6cc7cc(OC)ccc7n6c8ccccc8</chem>

No.	Structure
75	
76	

#### Example 4

##### HCV Replicon Cell Assay Protocol

- [0172] Cells containing hepatitis C virus (HCV) replicon were maintained in DMEM containing 10% fetal bovine serum (FBS), 0.25 mg per ml of G418, with appropriate supplements (media A).
- [0173] On day 1, replicon cell monolayer was treated with a trypsin:EDTA mixture, removed, and then media A was diluted into a final concentration of 100,000 cells per ml wit. 10,000 cells in 100 ul were plated into each well of a 96-well tissue culture plate, and cultured overnight in a tissue culture incubator at 37°C.

[0174] On day 2, compounds (in 100% DMSO) were serially diluted into DMEM containing 2% FBS, 0.5% DMSO, with appropriate supplements (media B). The final concentration of DMSO was maintained at 0.5% throughout the dilution series.

[0175] Media on the replicon cell monolayer was removed, and then media B containing various concentrations of compounds was added. Media B without any compound was added to other wells as no compound controls.

[0176] Cells were incubated with compound or 0.5% DMSO in media B for 48 hours in a tissue culture incubator at 37°C. At the end of the 48-hour incubation, the media was removed, and the replicon cell monolayer was washed once with PBS and stored at -80°C prior to RNA extraction.

[0177] Culture plates with treated replicon cell monolayers were thawed, and a fixed amount of another RNA virus, such as Bovine Viral Diarrhea Virus (BVDV) was added to cells in each well. RNA extraction reagents (such as reagents from RNeasy kits) were added to the cells immediately to avoid degradation of RNA. Total RNA was extracted according the instruction of manufacturer with modification to improve extraction efficiency and consistency. Finally, total cellular RNA, including HCV replicon RNA, was eluted and stored at -80°C until further processing.

[0178] A Taqman real-time RT-PCR quantification assay was set up with two sets of specific primers and probe. One was for HCV and the other was for BVDV. Total RNA extractants from treated HCV replicon cells was added to the PCR reactions for quantification of both HCV and BVDV RNA in the same PCR well. Experimental failure was flagged and rejected based on the level of BVDV RNA in

each well. The level of HCV RNA in each well was calculated according to a standard curve run in the same PCR plate. The percentage of inhibition or decrease of HCV RNA level due to compound treatment was calculated using the DMSO or no compound control as 0% of inhibition. The IC<sub>50</sub> (concentration at which 50% inhibition of HCV RNA level is observed) was calculated from the titration curve of any given compound.

Example 5

10 HCV Ki Assay Protocol

[0179] HPLC Microbore method for separation of 5AB substrate and products

Substrate:

NH<sub>2</sub>-Glu-Asp-Val-Val-(alpha)Abu-Cys-Ser-Met-Ser-Tyr-COOH

15 [0180] A stock solution of 20 mM 5AB (or concentration of your choice) was made in DMSO w/ 0.2M DTT. This was stored in aliquots at -20 C.

[0181] Buffer: 50 mM HEPES, pH 7.8; 20% glycerol; 100 mM NaCl

20 [0182] Total assay volume was 100  $\mu$ L

	X1 ( $\mu$ L)	conc. in assay
Buffer	86.5	see above
5 mM KK4A	0.5	25 $\mu$ M
1 M DTT	0.5	5 mM
DMSO or inhibitor	2.5	2.5% v/v
50 $\mu$ M tNS3	0.05	25 nM
250 $\mu$ M 5AB (initiate)	20	25 $\mu$ M

[0183] The buffer, KK4A, DTT, and tNS3 were combined; distributed 78  $\mu$ L each into wells of 96 well plate. This was incubated at 30 C for ~5-10 min.

[0184] 2.5  $\mu$ L of appropriate concentration of test compound was dissolved in DMSO (DMSO only for control) and added to each well. This was incubated at room temperature for 15 min.

- 5 [0185] Initiated reaction by addition of 20  $\mu$ L of 250  $\mu$ M 5AB substrate (25  $\mu$ M concentration is equivalent or slightly lower than the  $K_m$  for 5AB).

Incubated for 20 min at 30 C.

Terminated reaction by addition of 25  $\mu$ L of 10% TFA

- 10 Transferred 120  $\mu$ L aliquots to HPLC vials

[0186] Separated SMSY product from substrate and KK4A by the following method:

Microbore separation method:

**Instrumentation: Agilent 1100**

- 15 Degasser G1322A

Binary pump G1312A

Autosampler G1313A

Column thermostated chamber G1316A

Diode array detector G1315A

- 20 **Column:**

Phenomenex Jupiter; 5 micron C18; 300 angstroms; 150x2 mm; P/O 00F-4053-B0

Column thermostat: 40 C

Injection volume: 100  $\mu$ L

- 25 Solvent A = HPLC grade water + 0.1% TFA

Solvent B = HPLC grade acetonitrile + 0.1% TFA

Time (min)	%B	Flow (ml/min)	Max press.
0	5	0.2	400
12	60	0.2	400
13	100	0.2	400
16	100	0.2	400
17	5	0.2	400

Stop time: 17 min

Post-run time: 10 min.

[0187] Compounds of this invention have been tested in either the Example 4 and/or the Example 5 assays and have  
5 been shown to have HCV NS3-NS4A protease inhibition activity. Certain preferred compounds of this invention have comparable cell (Example 4) and enzymology (Example 5) data. Compounds 2, 7, 12, 13, 14, 18, 22, 24, 26, 34, 45, 50, 53, 54, 55, 57, 59, 61, 65, and 66, have  
10 comparable cell and enzyme data. More preferred compounds are 24, 45, 53, 54, 59, and 61 having comparable enzyme and cell data and both types of data falling within Category A.

[0188] Compounds of this invention were tested  
15 according to Example 5 (enzyme) and found to have  $K_i$  values of  $<0.1 \mu\text{M}$  (Category A);  $0.1-0.3 \mu\text{M}$  (Category B); and  $>0.3 \mu\text{M}$  (Category C) as follows.

Category A: 15, 19, 20, 24, 32, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 49, 51, 52, 53, 54, 56, 58, 59,  
20 61, 64, 67, and 69.

Category B: 1, 2, 3, 4, 5, 7, 8, 10, 12, 13, 14, 16, 18, 22, 23, 25, 30, 31, 33, 26, 34, 48, 50, 55, 57, 60, 65, 66, 70, 71, and 76.

Category C: 6, 9, 11, 17, 21, 27, 28, 29, 47 74, 75, and  
25 76.

[0189] Compounds of this invention were tested according to Example 4 (cell) and found to have  $\text{IC}_{50}$  values of  $<0.5 \mu\text{M}$  (Category A);  $0.5-1.0 \mu\text{M}$  (Category B); and  $>1.0 \mu\text{M}$  (Category C) as follows.

30 Category A: 7, 12, 13, 14, 15, 18, 19, 22, 24, 34, 35, 36, 38, 39, 40, 42, 43, 44, 45, 46, 47, 48, 50, 53, 54, 55, 56, 26, 57, 58, 59, 61, 64, 65, 66, and 67.

Category B: 1, 2, 3, 4, 5, 8, 10, 16, 20, and 70.

Category C: 6, 11, 41, 46, 47, 48, 49, 51, 60, and 71.

[0190] While we have described a number of embodiments of this invention, it is apparent that our basic examples may be altered to provide other embodiments which utilize the compounds and methods of this invention. Therefore, it will be appreciated that the scope of this invention is to be defined by the appended claims rather than by the specific embodiments that have been represented by way of example above. All cited documents are incorporated herein by reference.

10


15



## CLAIMS

We claim:

1. A compound of formula I:



# I

wherein:

Ar is a 5- to 10-membered aromatic ring having up to 4 heteroatoms selected from O, S, N(H), SO, and SO<sub>2</sub>, wherein 1 to 3 ring atoms are optionally and independently substituted with J;

$R^1$  and  $R^2$  are independently:

(C1-C12)-aliphatic-,

(C3-C10)-cycloalkyl- or -cycloalkenyl-,

15 [(C3-C10)-cycloalkyl- or -cycloalkenyl]-(C1-C12)-  
aliphatic-,

(C6-C10)-aryl- (C1-C12) aliphatic-,

(C6-C10)-heteroaryl- (C1-C12) aliphatic-,

wherein up to 3 aliphatic carbon atoms in R<sup>1</sup> and R<sup>2</sup> may be replaced by a heteroatom selected from O, N, S, SO, or SO<sub>2</sub> in a chemically stable arrangement;

wherein each of R<sup>1</sup> and R<sup>2</sup> is independently and optionally substituted with up to 3 substituents independently selected from J;

25

R<sup>3</sup> and R<sup>3'</sup> are independently hydrogen or (C1-C12)-  
aliphatic, wherein any hydrogen is optionally replaced  
with halogen; wherein any terminal carbon atom of R<sup>3</sup> is  
optionally substituted with sulfhydryl or hydroxy; or  
R<sup>3</sup> is phenyl or -CH<sub>2</sub>phenyl, wherein said phenyl group


is optionally substituted with up to 3 substituents independently selected from J; or  
R<sup>3</sup> and R<sup>3'</sup> together with the atom to which they are bound is a 3- to 6-membered ring having up to 2 heteroatoms selected from N, NH, O, SO, and SO<sub>2</sub>; wherein the ring  
5 has up to 2 substituents selected independently from J;

R<sup>4</sup> and R<sup>4'</sup> are independently:  
hydrogen-,  
10 (C1-C12)-aliphatic-,  
(C3-C10)-cycloalkyl- or -cycloalkenyl-,  
(C3-C10)-cycloalkyl-(C1-C12)-aliphatic-,  
(C6-C10)-aryl-,  
(C3-C10)-heterocyclyl-; or  
15 (C5-C10)-heteroaryl-;

wherein up to two aliphatic carbon atoms in R<sup>4</sup> and R<sup>4'</sup> may be replaced by a heteroatom selected from O, N, S, SO, and SO<sub>2</sub>;

wherein each of R<sup>4</sup> and R<sup>4'</sup> is independently and  
20 optionally substituted with up to 3 substituents independently selected from J;

W is:



25 wherein each R<sub>6</sub> is independently:  
hydrogen-,

(C1-C12)-aliphatic-,  
(C6-C10)-aryl-,  
(C6-C10)-aryl-(C1-C12)aliphatic-,  
(C3-C10)-cycloalkyl- or cycloalkenyl-,  
5 [(C3-C10)-cycloalkyl- or cycloalkenyl]-(C1-C12)-  
aliphatic-,  
(C3-C10)-heterocyclyl-,  
(C3-C10)-heterocyclyl-(C1-C12)-aliphatic-,  
(C5-C10)heteroaryl-, or  
10 (C5-C10)heteroaryl-(C1-C12)-aliphatic-, or  
two R<sub>6</sub> groups, which are bound to the same  
nitrogen atom, form together with that nitrogen  
atom, a (C3-C10)-heterocyclic ring;  
wherein R<sub>6</sub> is optionally substituted with up to 3  
15 J substituents;  
wherein each R<sub>8</sub> is independently -OR'; or the R<sub>8</sub>  
groups together with the boron atom, is a (C3-C10)-  
membered heterocyclic ring having in addition to the  
boron up to 3 additional heteroatoms selected from N,  
20 NH, O, SO, and SO<sub>2</sub>;

T is:

(C1-C12)-aliphatic-;  
(C6-C10)-aryl-,  
25 (C6-C10)-aryl-(C1-C12)aliphatic-,  
(C5-C10)heteroaryl-, or  
(C5-C10)heteroaryl-(C1-C12)-aliphatic-;  
wherein each T is optionally substituted with up to  
3 J substituents;  
30 J is halogen, -OR', -NO<sub>2</sub>, -CN, -CF<sub>3</sub>, -OCF<sub>3</sub>, -R', oxo,  
thioxo, 1,2-methylenedioxy, 1,2-ethylenedioxy, =N(R'),  
=N(OR'), -N(R')<sub>2</sub>, -SR', -SOR', -SO<sub>2</sub>R', -SO<sub>2</sub>N(R')<sub>2</sub>,  
-SO<sub>3</sub>R', -C(O)R', -C(O)C(O)R', -C(O)CH<sub>2</sub>C(O)R', -C(S)R',


-C(S)OR', -C(O)OR', -C(O)C(O)OR', -C(O)C(O)N(R')<sub>2</sub>,  
 -OC(O)R', -C(O)N(R')<sub>2</sub>, -OC(O)N(R')<sub>2</sub>, -C(S)N(R')<sub>2</sub>,  
 -(CH<sub>2</sub>)<sub>0-2</sub>NHC(O)R', -N(R')N(R')COR', -N(R')N(R')C(O)OR',  
 -N(R')N(R')CON(R')<sub>2</sub>, -N(R')SO<sub>2</sub>R', -N(R')SO<sub>2</sub>N(R')<sub>2</sub>,  
 5 -N(R')C(O)OR', -N(R')C(O)R', -N(R')C(S)R',  
 -N(R')C(O)N(R')<sub>2</sub>, -N(R')C(S)N(R')<sub>2</sub>, -N(COR')COR',  
 -N(OR')R', -C(=NH)N(R')<sub>2</sub>, -C(O)N(OR')R', -C(=NOR')R',  
 -OP(O)(OR')<sub>2</sub>, -P(O)(R')<sub>2</sub>, -P(O)(OR')<sub>2</sub>, or -P(O)(H)(OR');

10 R' is:

hydrogen-,  
 (C1-C12)-aliphatic-,  
 (C3-C10)-cycloalkyl- or -cycloalkenyl-,  
 [(C3-C10)-cycloalkyl or -cycloalkenyl]-(C1-C12)-  
 15 aliphatic-,  
 (C6-C10)-aryl-,  
 (C6-C10)-aryl-(C1-C12)aliphatic-,  
 (C3-C10)-heterocyclyl-,  
 (C6-C10)-heterocyclyl-(C1-C12)aliphatic-,  
 20 (C5-C10)-heteroaryl-, or  
 (C5-C10)-heteroaryl-(C1-C12)-aliphatic-;  
 wherein R' is optionally substituted with up to 3  
 J groups;

wherein two R' groups bound to the same atom  
 25 form a 3- to 10-membered aromatic or non-aromatic  
 ring having up to 3 heteroatoms independently  
 selected from N, O, S, SO, or SO<sub>2</sub>, wherein said  
 ring is optionally fused to a (C6-C10)aryl, (C5-  
 C10)heteroaryl, (C3-C10)cycloalkyl, or a (C3-  
 30 C10)heterocyclyl, wherein any ring has up to 3  
 substituents selected independently from J.

2. A compound of formula I:



I

wherein:

n is 0 or 1;

- 5 Ar is a 5- to 10-membered aromatic ring having up to 4 heteroatoms selected from O, S, N(H), SO, and SO<sub>2</sub>, wherein 1 to 3 ring atoms are optionally and independently substituted with J;

R<sup>1</sup>, R<sup>2</sup>, R<sup>12</sup>, and R<sup>13</sup> are independently:

- 10 (C1-C12)-aliphatic-,  
 (C3-C10)-cycloalkyl- or -cycloalkenyl-,  
 [(C3-C10)-cycloalkyl- or -cycloalkenyl]-(C1-C12)-  
 aliphatic-,  
 (C6-C10)-aryl-(C1-C12)aliphatic-,  
 15 (C6-C10)-heteroaryl-(C1-C12)aliphatic-,

wherein up to 3 aliphatic carbon atoms in R<sup>1</sup> and R<sup>2</sup> may be replaced by a heteroatom selected from O, N, NH, S, SO, or SO<sub>2</sub> in a chemically stable arrangement;

- wherein each of R<sup>1</sup> and R<sup>2</sup> is independently and  
 20 optionally substituted at each substitutable position with up to 3 substituents independently selected from J;

R<sup>3</sup> and R<sup>3'</sup> are independently hydrogen or (C1-C12)-  
 aliphatic, wherein any hydrogen is optionally replaced  
 25 with halogen; wherein any terminal carbon atom of R<sup>3</sup> is optionally substituted with sulfhydryl or hydroxy; or R<sup>3</sup> is phenyl or -CH<sub>2</sub>phenyl, wherein said phenyl group is optionally substituted with up to 3 substituents independently selected from J; or

- 30 R<sup>3</sup> and R<sup>3'</sup> together with the atom to which they are bound is a 3- to 6-membered ring having up to 2 heteroatoms

selected from N, NH, O, SO, and SO<sub>2</sub>; wherein the ring has up to 2 substituents selected independently from J;


R<sup>4</sup> and R<sup>4'</sup> are independently:

- 5 hydrogen-,  
 (C1-C12)-aliphatic-,  
 (C3-C10)-cycloalkyl- or -cycloalkenyl-,  
 (C3-C10)-cycloalkyl-(C1-C12)-aliphatic-,  
 (C6-C10)-aryl-,  
 10 (C3-C10)-heterocyclcyl-; or  
 (C5-C10)-heteroaryl-;

wherein up to two aliphatic carbon atoms in R<sup>4</sup> and R<sup>4'</sup> may be replaced by a heteroatom selected from O, N, S, SO, and SO<sub>2</sub>;

- 15 wherein each of R<sup>4</sup> and R<sup>4'</sup> is independently and optionally substituted with up to 3 substituents independently selected from J;

W is:



20

wherein

Y is -CO<sub>2</sub>H, a derivative of -CO<sub>2</sub>H, or a bioisostere of -CO<sub>2</sub>H;

each R<sub>6</sub> is independently:

- 25 hydrogen-,

(C1-C12)-aliphatic-,  
(C6-C10)-aryl-,  
(C6-C10)-aryl-(C1-C12)aliphatic-,  
(C3-C10)-cycloalkyl- or cycloalkenyl-,  
5 [(C3-C10)-cycloalkyl- or cycloalkenyl]-(C1-C12)-  
aliphatic-,  
(C3-C10)-heterocyclyl-,  
(C3-C10)-heterocyclyl-(C1-C12)-aliphatic-,  
(C5-C10)heteroaryl-, or  
10 (C5-C10)heteroaryl-(C1-C12)-aliphatic-, or  
two R<sub>6</sub> groups, which are bound to the same  
nitrogen atom, form together with that nitrogen  
atom, a (C3-C10)-heterocyclic ring;  
wherein R<sub>6</sub> is optionally substituted with up to 3  
15 J substituents;  
wherein each R<sub>8</sub> is independently -OR'; or the R<sub>8</sub>  
groups together with the boron atom, is a (C3-C10)-  
membered heterocyclic ring having in addition to the  
boron up to 3 additional heteroatoms selected from N,  
20 NH, O, SO, and SO<sub>2</sub>;

T is:  
(C1-C12)-aliphatic-;  
(C6-C10)-aryl-,  
25 (C6-C10)-aryl-(C1-C12)aliphatic-,  
(C5-C10)heteroaryl-, or  
(C5-C10)heteroaryl-(C1-C12)-aliphatic-;  
wherein each T is optionally substituted with up to  
3 J substituents; and  
30 wherein up to 3 aliphatic carbon atoms in T may be  
replaced by a heteroatom selected from O, N, NH, S, SO,  
or SO<sub>2</sub> in a chemically stable arrangement;  
provided that if T is pyrrole, the pyrrole is not  
substituted at the 3-position with J, with J being

-C(O)R', -C(O)C(O)R', -C(O)CH<sub>2</sub>C(O)R', -C(S)R', -C(S)OR',  
 -C(O)OR', -C(O)C(O)OR', -C(O)C(O)N(R')<sub>2</sub>, -C(O)N(R')<sub>2</sub>,  
 -C(S)N(R')<sub>2</sub>, -C(=NH)N(R')<sub>2</sub>, -C(O)N(OR')R', -C(=NOR')R';

J is halogen, -OR', -NO<sub>2</sub>, -CN, -CF<sub>3</sub>, -OCF<sub>3</sub>, -R', oxo,  
 thioxo, 1,2-methylenedioxy, 1,2-ethylenedioxy, =N(R'),  
 =N(OR'), -N(R')<sub>2</sub>, -SR', -SOR', -SO<sub>2</sub>R', -SO<sub>2</sub>N(R')<sub>2</sub>,  
 -SO<sub>3</sub>R', -C(O)R', -C(O)C(O)R', -C(O)CH<sub>2</sub>C(O)R', -C(S)R',  
 -C(S)OR', -C(O)OR', -C(O)C(O)OR', -C(O)C(O)N(R')<sub>2</sub>,  
 -OC(O)R', -C(O)N(R')<sub>2</sub>, -OC(O)N(R')<sub>2</sub>, -C(S)N(R')<sub>2</sub>,  
 -(CH<sub>2</sub>)<sub>0-2</sub>NHC(O)R', -N(R')N(R')COR', -N(R')N(R')C(O)OR',  
 -N(R')N(R')CON(R')<sub>2</sub>, -N(R')SO<sub>2</sub>R', -N(R')SO<sub>2</sub>N(R')<sub>2</sub>,  
 -N(R')C(O)OR', -N(R')C(O)R', -N(R')C(S)R',  
 -N(R')C(O)N(R')<sub>2</sub>, -N(R')C(S)N(R')<sub>2</sub>, -N(COR')COR',  
 -N(OR')R', -C(=NH)N(R')<sub>2</sub>, -C(O)N(OR')R', -C(=NOR')R',  
 -OP(O)(OR')<sub>2</sub>, -P(O)(R')<sub>2</sub>, -P(O)(OR')<sub>2</sub>, or -P(O)(H)(OR');

R' is:

hydrogen-,  
 (C1-C12)-aliphatic-,  
 (C3-C10)-cycloalkyl- or -cycloalkenyl-,  
 [(C3-C10)-cycloalkyl or -cycloalkenyl]-(C1-C12)-  
 aliphatic-,  
 (C6-C10)-aryl-,  
 (C6-C10)-aryl-(C1-C12)aliphatic-,  
 (C3-C10)-heterocyclyl-,  
 (C6-C10)-heterocyclyl-(C1-C12)aliphatic-,  
 (C5-C10)-heteroaryl-, or  
 (C5-C10)-heteroaryl-(C1-C12)-aliphatic-;

wherein R' is optionally substituted with up to 3

J groups;


wherein two R' groups bound to the same atom form a 3- to  
 10-membered aromatic or non-aromatic ring having up to 3  
 heteroatoms independently selected from N, O, S, SO, or




SO<sub>2</sub>, wherein said ring is optionally fused to a (C6-C10)aryl, (C5-C10)heteroaryl, (C3-C10)cycloalkyl, or a (C3-C10)heterocyclyl, wherein any ring has up to 3 substituents selected independently from J.

3. The compound according to claim 1 or claim 2, wherein Ar is phenyl, pyridyl, quinolinyl, pyrimidinyl, or naphthyl, wherein each group is optionally substituted with 1, 2, or 3 J groups.

4. The compound according to claim 1 or claim 2, wherein Ar is



5. The compound according to claim 1 or claim 2, wherein Ar is:




6. The compound according to claim 1 or claim 2, wherein Ar is a 6 or a 10-membered aromatic ring having 0, 1, or 2 nitrogen heteroatoms, wherein 1, 2, or 3 ring atoms are optionally and independently substituted with J.

7. The compound according to any one of claims 1-6, wherein each J group on Ar is independently OR', NO<sub>2</sub>, CN, CF<sub>3</sub>, OCF<sub>3</sub>, R', COR', C(O)OR', C(O)N(R')<sub>2</sub>, SO<sub>2</sub>R', SO<sub>2</sub>N(R')<sub>2</sub>, 1,2-methylenedioxy, 1,2-ethylene dioxy, or NR'C(O)OR', NR'SO<sub>2</sub>R'.

8. The compound according to any one of claims 1-6, wherein each J group on Ar is independently OR', halogen, CN, CF<sub>3</sub>, R', or COR'.


9. The compound according to any one of claims 1-6, wherein each J group on Ar is independently halo, trifluoromethyl, methyl, or NO<sub>2</sub>.

10. The compound according to any one of claims 1-9, wherein W is:




wherein the NR<sup>6</sup>R<sup>6</sup> is -NH-(C1-C6 aliphatic), -NH-(C3-C6 cycloalkyl), -NH-CH(CH<sub>3</sub>)-aryl, or -NH-CH(CH<sub>3</sub>)-heteroaryl, wherein said aryl or said heteroaryl is optionally substituted with up to 3 halogens.


11. The compound according to any one of claims 1-10, wherein in the W, the  $\text{NR}^6\text{R}^6$  is:




12. The compound according to any one of claims 1-10, wherein in the W, the  $\text{NR}^6\text{R}^6$  is:




13. The compound according to any one of claims 1-10, wherein in the W, the  $\text{NR}^6\text{R}^6$  is:




14. The compound according to any one of claims 1-10, wherein in the W, the  $\text{NR}^6\text{R}^6$  is:




15. The compound according to any one of claims 1-14, wherein R<sup>3'</sup> is hydrogen and R<sup>3</sup> is:




16. The compound according to any one of claims 1-14, wherein R<sup>3'</sup> is hydrogen and R<sup>3</sup> is:




17. The compound according to any one of claims 1-16, wherein  $R^2$  is:




18. The compound according to claim 17, wherein R<sup>2</sup> is:




19. The compound according to claim 18, wherein  $R^2$  is:



20. The compound according to any one of claims 1-19, wherein  $R^1$  is:




21. The compound according to claim 20, wherein  $R^1$  is:



22. The compound according to claim 21, wherein  $R^1$  is cyclohexyl.


23. The compound according to any one of claims 1-22, wherein T is (C6-C10)-aryl- or (C5-C10)heteroaryl-, wherein each T is optionally substituted with 1, 2, or 3 J substituents.

24. The compound according to any one of claims 1-22, wherein T is:









wherein each T group is optionally fused to a 5-membered or a 6-membered aryl or heteroaryl group.

- 5            25. The compound according to claim 24, wherein T is:









- 10            26. The compound according to claim 1 wherein, the compound is:



No.	Structure
1	 <chem>CC1(C)C(=O)N(C1C(=O)N(C(C2CCCCC2)C(=O)N3C(=O)c4ccncc4C3=O)C(=O)N5C(=O)C(C5)OC6=CC(=CC(=C6)F)F)C(=O)N[C@H](C)C(=O)N7C(=O)C8CC8C7=O</chem>
2	 <chem>CC1(C)C(=O)N(C1C(=O)N(C(C2CCCCC2)C(=O)N3C(=O)c4ccncc4C3=O)C(=O)N[C@H](C)C(=O)N5C(=O)C(C5)OC6=CC(=CC(=C6)Cl)Cl)C(=O)N[C@H](C)C(=O)N7C(=O)C8CC8C7=O</chem>
3	 <chem>CC1(C)C(=O)N(C1C(=O)N(C(C2CCCCC2)C(=O)N3C(=O)c4ccncc4C3=O)C(=O)N[C@H](C)C(=O)N5C(=O)C(C5)OC6=CC(=CC(=C6)Cl)Cl)C(=O)N[C@H](C)C(=O)N7C(=O)C8CC8C7=O</chem>




No.	Structure
4	
5	
6	








No.	Structure
7	
8	
9	




No.	Structure
10	 <chem>CC[C@H](NC(=O)C1CCN1C(=O)N[C@@H](C(C)(C)C)C(=O)N[C@@H](C2CCCCC2)C(=O)c3ccncc3)C(=O)N[C@@H](COc4cc(F)c(F)cc4)C(=O)N[C@@H](CC)C(=O)N5CC5</chem>
11	 <chem>CC[C@H](NC(=O)C1CCN1C(=O)N[C@@H](C(C)(C)C)C(=O)N[C@@H](C2CCCCC2)C(=O)c3ccncc3)C(=O)N[C@@H](COc4cc(C(F)(F)F)c(C(F)(F)F)cc4)C(=O)N[C@@H](CC)C(=O)N5CC5</chem>
12	 <chem>CC[C@H](NC(=O)C1CCN1C(=O)N[C@@H](C(C)(C)C)C(=O)N[C@@H](C2CCCCC2)C(=O)c3ccncc3)C(=O)N[C@@H](COc4cc(Cl)c(Cl)cc4)C(=O)N[C@@H](CC)C(=O)N5CC5</chem>



No.	Structure
13	 <chem>CC[C@H](NC(=O)N1CC[C@H](C1)C2=CC=C(C=C2)OC(F)(F)F)C(=O)N[C@@H](C(C)(C)C)C(=O)N[C@@H](C3CCCCC3)C(=O)c4ccncc4</chem>
14	 <chem>CC[C@H](NC(=O)N1CC[C@H](C1)C2=CC=C(C=C2)OC(F)(F)F)C(=O)N[C@@H](C(C)(C)C)C(=O)N[C@@H](C3CCCCC3)C(=O)c4ccncc4</chem>

No.	Structure
15	 <chem>CC[C@H](NC(=O)C1CCN(C1)C(=O)N[C@@H](C(C)(C)C)C(=O)N[C@@H](C2CCCCC2)C(=O)Nc3ccncc3)Oc4ccc(cc4Oc5ccccc5c6ccccc6)C(=O)N7CC8C7C8</chem>
16	 <chem>CC[C@H](NC(=O)C1CCN(C1)C(=O)N[C@@H](C(C)(C)C)C(=O)N[C@@H](C2CCCCC2)C(=O)Nc3ccncc3)Oc4ccc(cc4OC(F)(F)F)c5ccccc5</chem>
17	 <chem>CC[C@H](NC(=O)C1CCN(C1)C(=O)N[C@@H](C(C)(C)C)C(=O)N[C@@H](C2CCCCC2)C(=O)Nc3ccncc3)Oc4ccccc4Oc5ccccc5</chem>




No.	Structure
18	
19	
20	




No.	Structure
21	 <p>Chemical structure 21 is a complex molecule. It features a pyridine ring connected via an amide bond to a cyclohexyl group. This is further linked to a chiral center with a tert-butyl group and a pyrrolidine ring. The pyrrolidine ring is connected to a chiral center with an ethyl group and a cyclopropyl group.</p>
22	 <p>Chemical structure 22 is a complex molecule, similar to structure 21, but with a trifluoromethyl group on the phenyl ring of the pyrrolidine moiety.</p>

No.	Structure
23	 <chem>COc1cc(OC)ccc1O[C@@H]1CC[C@H](C(=O)N[C@@H](CC)C(=O)NC2CC2)N1C(=O)N[C@@H](C(C)(C)C)C(=O)N[C@@H](C1CCCCC1)C(=O)c1ccncc1</chem>
24	 <chem>COc1cc(OC)cc(OC)c1O[C@@H]1CC[C@H](C(=O)N[C@@H](CC)C(=O)NC2CC2)N1C(=O)N[C@@H](C(C)(C)C)C(=O)N[C@@H](C1CCCCC1)C(=O)c1ccncc1</chem>
25	 <chem>Clc1ccncc1O[C@@H]1CC[C@H](C(=O)N[C@@H](CC)C(=O)NC2CC2)N1C(=O)N[C@@H](C(C)(C)C)C(=O)N[C@@H](C1CCCCC1)C(=O)c1ccncc1</chem>



No.	Structure
26	 <chem>CC[C@H](C(=O)N[C@@H](Cc1ccc(cc1)[N+](=O)[O-])C2CCCN2C(=O)N[C@@H](C(C)(C)C)C(=O)N[C@@H](C3CCCCC3)C(=O)c4ccncc4)C(=O)N[C@@H](C5CC5)C(=O)C</chem>
27	 <chem>CC[C@H](C(=O)N[C@@H](Cc1ccc(cc1)C(F)(F)F)C2CCCN2C(=O)N[C@@H](C(C)(C)C)C(=O)N[C@@H](C3CCCCC3)C(=O)c4ccncc4)C(=O)N[C@@H](C5CC5)C(=O)C</chem>









No.	Structure
28	
29	
30	




No.	Structure
31	 <chem>CC(C)(C)C(=O)N[C@@H](Cc1cccnc1)C(=O)N[C@@H](Cc2ccccc2)c3ccccc3</chem>
32	 <chem>CC(C)(C)C(=O)N[C@@H](Cc1cccnc1)C(=O)N[C@@H](Cc2ccccc2)c3ccccc3</chem>
33	 <chem>CC(C)(C)C(=O)N[C@@H](Cc1cccnc1)C(=O)N[C@@H](Cc2ccccc2)c3ccccc3</chem>



No.	Structure
34	 <chem>CN1CC[C@H](C1)C(=O)N[C@@H](CCC)C(=O)N[C@@H](C)C(=O)N[C@@H](C(C)(C)C)C(=O)N[C@@H](C2CCCCC2)C(=O)Nc3ccncc3Oc4ccccc4</chem>
35	 <chem>CN1CC[C@H](C1)C(=O)N[C@@H](CCC)C(=O)N[C@@H](C)C(=O)N[C@@H](C(C)(C)C)C(=O)N[C@@H](C2CCCCC2)C(=O)Nc3ccncc3Oc4ccccc4</chem>
36	 <chem>CN1CC[C@H](C1)C(=O)N[C@@H](CCC)C(=O)N[C@@H](C)C(=O)N[C@@H](C(C)(C)C)C(=O)N[C@@H](C2CCCCC2)C(=O)Nc3ccncc3Oc4c5ccccc5cc4</chem>

No.	Structure
37	 <p>Chemical structure 37 is a complex molecule. It features a naphthalen-1-yloxy group attached to a pyrrolidine ring. The pyrrolidine ring is part of a chain containing a quaternary carbon with two methyl groups, a cyclohexyl group, a pyrazole ring, and a cyclopropyl group.</p>
38	 <p>Chemical structure 38 is a complex molecule. It features a 6-methoxynaphthalen-1-yloxy group attached to a pyrrolidine ring. The pyrrolidine ring is part of a chain containing a quaternary carbon with two methyl groups, a cyclohexyl group, a pyridine ring, and a cyclopropyl group.</p>



No.	Structure
39	
40	
41	




No.	Structure
42	
43	
44	




No.	Structure
45	
46	
47	




No.	Structure
48	
49	








No.	Structure
50	
51	




No.	Structure
52	 <chem>COc1ccc(O[C@@H]2CC[C@H](C(=O)N[C@@H](CC)C(=O)N[C@@H](C3CCCCC3)C(=O)Nc4ccncc4)C2)c(C)C</chem>
53	 <chem>Clc1cc(Cl)cc(OC[C@@H]2CC[C@H](C(=O)N[C@@H](CC)C(=O)N[C@@H](C3CCCCC3)C(=O)Nc4ccncc4)C2)c1</chem>
54	 <chem>Fc1cc(F)cc(OC[C@@H]2CC[C@H](C(=O)N[C@@H](CC)C(=O)N[C@@H](C3CCCCC3)C(=O)Nc4ccncc4)C2)c1</chem>



No.	Structure
55	 <chem>CC[C@H](C)[C@@H](C1CCN(C1)C(=O)N[C@@H](C)[C@H](C2=CC=CC=C2F)OCC3=CC=C(C=C3)F)C(=O)N[C@@H](C)[C@H](C4=CC=CC=N4)C(=O)N[C@@H](C5CCCCC5)C(=O)N</chem>
56	 <chem>CC[C@H](C)[C@@H](C1CCN(C1)C(=O)N[C@@H](C)[C@H](C2=CC=C(C=C2)Cl)OCC3=CC=CC=C3)C(=O)N[C@@H](C)[C@H](C4=CC=CC=N4)C(=O)N[C@@H](C5CCCCC5)C(=O)N</chem>
57	 <chem>CC[C@H](C)[C@@H](C1CCN(C1)C(=O)N[C@@H](C)[C@H](C2=CC=C(C=C2)[N+](=O)[O-])OCC3=CC=CC=C3)C(=O)N[C@@H](C)[C@H](C4=CC=CC=N4)C(=O)N[C@@H](C5CCCCC5)C(=O)N</chem>

No.	Structure
58	
59	
60	



No.	Structure
61	 <chem>CC(C)(C)C(NC(=O)C1CCCCC1C(=O)Nc2cccnc2)C(=O)N[C@@H](CC)C(=O)NCC(=O)N[C@@H](CC)C(=O)NCC(=O)Nc3ccc(OC4CCCCC4C(=O)Nc5cccnc5)cc3</chem>
62	 <chem>CC(C)(C)C(NC(=O)C1CCCCC1C(=O)Nc2ccc(Cl)cc2)C(=O)N[C@@H](CC)C(=O)N[C@@H](CC)C(=O)NCC(=O)N[C@@H](CC)C(=O)Nc3ccc(OC4CCCCC4C(=O)Nc5cccnc5)cc3</chem>



No.	Structure
63	 <chem>CC1(C)C(C2=CC=CC=C2OC3CC[C@H](C(=O)NCC(=O)NCC(=O)NCC4CC4)N3C(=O)N[C@@H](C5CCCCC5)C(=O)N6C=CC=CC=N6)C1</chem>
64	 <chem>CC1(C)C(C2=CC(=CC=C2)OC3CC[C@H](C(=O)NCC(=O)NCC(=O)NCC4CC4)N3C(=O)N[C@@H](C5CCCCC5)C(=O)N6C=CC=CC=N6)C1</chem>
65	 <chem>N#CC1=CC=C(OC2CC[C@H](C(=O)NCC(=O)NCC(=O)NCC3CC3)N2)C=C1</chem>



No.	Structure
66	 <chem>CC[C@H](C)[C@@H](C1CCN(C1)C(=O)N[C@@H](C)[C@H](C2CCCCC2)C(=O)Nc3ccncc3)C(=O)N[C@@H](C)[C@H](C4CCCCC4)C(=O)Nc5ccncc5</chem>
67	 <chem>CC[C@H](C)[C@@H](C1CCN(C1)C(=O)N[C@@H](C)[C@H](C2CCCCC2)C(=O)Nc3ccncc3)C(=O)N[C@@H](C)[C@H](C4CCCCC4)C(=O)Nc5ccc6ncncc6c5C(F)(F)F</chem>
68	 <chem>CC[C@H](C)[C@@H](C1CCN(C1)C(=O)N[C@@H](C)[C@H](C2CCCCC2)C(=O)Nc3ccncc3)C(=O)N[C@@H](C)[C@H](C4CCCCC4)C(=O)Nc5cc(F)c(F)cc5</chem>

No.	Structure
69	 <p>Chemical structure 69 is a complex molecule. It features a central cyclohexyl group connected to a 4-chlorophenyl group via a 4-ethyl-1H-imidazole-5-carboxamide moiety. The structure includes a 4-ethyl-1H-imidazole-5-carboxamide moiety, a 4-chlorophenyl group, and a cyclohexyl group.</p>
70	 <p>Chemical structure 70 is a complex molecule. It features a central cyclohexyl group connected to a 4-ethyl-1H-imidazole-5-carboxamide moiety. The structure includes a 4-ethyl-1H-imidazole-5-carboxamide moiety, a 4-chlorophenyl group, and a cyclohexyl group.</p>



No.	Structure
71	 <p>Chemical structure 71 is a complex molecule. It features a pyridine ring connected via an amide bond to a cyclohexyl group. This is further linked to a central chiral center (a carbon atom bonded to a methyl group, a tert-butyl group, and a hydrogen atom). The central chiral center is also connected to a pyrrolidine ring, which is linked via an ether bond to a 4-(piperidin-1-yl)benzoyl group. The pyrrolidine ring is also connected to a chain of amide bonds, including a cyclopropyl group and an ethyl group.</p>
72	 <p>Chemical structure 72 is a complex molecule, similar to structure 71, but with a 4-chloro-2-pyridyl group instead of the 4-(piperidin-1-yl)benzoyl group. The rest of the molecule, including the pyridine ring, cyclohexyl group, central chiral center, and the chain of amide bonds, is identical to structure 71.</p>

No.	Structure
73	 <chem>CCOC1=CC=C2C(=C1)N=C(C=C2)C3=CC=CC=C3[C@@H]3CC[C@H](C(=O)N[C@@H](CC)C(=O)N[C@@H](C4CCCCC4)C(=O)Nc5ccncc5)[C@H]3N</chem>
74	 <chem>CCOC1=CC=C2C(=C1)N=C(C=C2)C(F)(F)F[C@H]3CC[C@H](C(=O)N[C@@H](CC)C(=O)N[C@@H](C4CCCCC4)C(=O)Nc5ccncc5)[C@H]3N</chem>

No.	Structure
75	
76	

27. A composition comprising a compound according to any one of claims 1-26 or a pharmaceutically acceptable salt thereof, in an amount effective to  
5 inhibit a serine protease; and a acceptable carrier, adjuvant or vehicle.

28. The composition according to claim 27, wherein said composition is formulated for administration to a  
10 patient.

29. The composition according to claim 28, wherein said composition comprises an additional agent selected

from an immunomodulatory agent; an antiviral agent; a second inhibitor of HCV protease; an inhibitor of another target in the HCV life cycle; a cytochrome P-450 inhibitor; or combinations thereof.

30. The composition according to claim 29, wherein said immunomodulatory agent is  $\alpha$ -,  $\beta$ -, or  $\gamma$ -interferon or thymosin; the antiviral agent is ribavirin, amantadine, or telbivudine; or the inhibitor of another target in the HCV life cycle is an inhibitor of HCV helicase, polymerase, or metalloprotease.

31. The composition according to claim 30, wherein said cytochrome P-450 inhibitor is ritonavir.

32. A method of inhibiting the activity of a serine protease comprising the step of contacting said serine protease with a compound according to any one of claims 1-26.

33. The method according to claim 32, wherein said protease is an HCV NS3 protease.

34. A method of treating an HCV infection in a patient comprising the step of administering to said patient a composition according to claim 27.

35. The method according to claim 34, comprising the additional step of administering to said patient an additional agent selected from an immunomodulatory agent; an antiviral agent; a second inhibitor of HCV protease; an inhibitor of another target in the HCV life cycle; or combinations thereof; wherein said additional agent is

administered to said patient as part of said composition according to claim 30 or as a separate dosage form.


36. The method according to claim 35, wherein said immunomodulatory agent is  $\alpha$ -,  $\beta$ -, or  $\gamma$ -interferon or thymosin; said antiviral agent is ribavarin or amantadine; or said inhibitor of another target in the HCV life cycle is an inhibitor of HCV helicase, polymerase, or metalloprotease.

37. A method of eliminating or reducing HCV contamination of a biological sample or medical or laboratory equipment, comprising the step of contacting said biological sample or medical or laboratory equipment with a compound according to any one of claims 1-26.

38. The method according to claim 37, wherein said sample or equipment is selected from a body fluid, biological tissue, a surgical instrument, a surgical garment, a laboratory instrument, a laboratory garment, a blood or other body fluid collection apparatus; a blood or other bodily fluid storage material.

39. The method according to claim 38, wherein said body fluid is blood.


40. A process for preparing a compound of formula I, as defined in any one of claims 1-26, comprising the step of: reacting a compound of formula II in the presence of a compound of formula III to provide a compound of formula IV:



II

ArX

III



IV

wherein:

R<sup>10</sup> is an amine protecting group, a P3- residue of an HCV protease inhibitor described herein, or a P4-P3- residue of an HCV protease inhibitor as described herein, and wherein the P3 and the P4-P3 residues are optionally protected the an amino-terminal capping group;

R<sup>11</sup> is a carboxy protecting group or a P1 residue of an HCV protease inhibitor described herein, wherein the P1 residue is optionally substituted at the terminal carboxy position with a carboxy protecting group or with W.